



Adv Biomed Res. 2017; 6: 49.

PMCID: PMC5433655

Published online 2017 Apr 25. doi: [10.4103/2277-9175.205191](https://doi.org/10.4103/2277-9175.205191)

## Phenotypic and Molecular Identification of Nocardia in Brain Abscess

[Mehdi Fatahi-Bafghi](#)

From the Department of Microbiology, Faculty of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

**Address for correspondence:** Dr. Mehdi Fatahi-Bafghi, Department of Microbiology, Faculty of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran. E-mail: [mehdifatahi@ssu.ac.ir](mailto:mehdifatahi@ssu.ac.ir)

Received 2015 Dec; Accepted 2016 Jun.

[Copyright](#) : © 2017 Advanced Biomedical Research

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Sir,

Shirani *et al.* recently reported an article entitled, “Nocardial brain abscess in a patient with pulmonary alveolar proteinosis” (DOI: 10.4103/2277-9175.164004).[1] The genus *Nocardia* is a Gram-positive aerobic, partially acid-fast, and filamentous bacterium that causes nocardial infections (nocardiosis) in human. The genus *Nocardia* is normal microflora in the environment such as soil and dust.[2] Nocardiosis treatment is different in various species, and some species are resistant to different antibiotics such as trimethoprim-sulfamethoxazole, carbapenem, and aminoglycosides, as well as accurate identification in species level is important.

Attention to comments and questions:

1. Isolation and characterization of morphology of colony, aerial hyphae, Gram stain, partially acid-fast, acid-fast, growth in lysozyme broth and molecular methods are important for the genus *Nocardia* confirmation.[2,3,4] Moreover, other aerobic actinomycetes such as *Gordonia* spp., *Rhodococcus* spp., and *Tsukamurella* spp. are Gram-positive, may be filamentous and colonial morphology are similar to each other. *Gordonia* spp., *Rhodococcus* spp., *Tsukamurella* spp., and *Nocardia* spp. are acid-fast in under certain conditions (bacteria listed are partially acid-fast and normally are not positive for acid-fast staining)[2,3,4,5,6,7,8,9,10,11,12] while authors suggested that branching, filamentous, Gram-positive, and acid-fast positive elements are the genus *Nocardia*
2. Authors reported *Nocardia asteroides* identification with culture and staining.[1] Phenotypic methods such as hydrolysis of amino acids, production of nitrate reductase, gelatinase, and urease, producing acid from carbohydrates, and growth at 45°C[2] are used in species level identification. I have two questions for authors:
  - i. The genus *Nocardia* has some of complex groups such as *N. asteroides* complex (*N. asteroides*, *Nocardia cyriacigeorgica*, *Nocardia farcinica*, etc.,) and *Nocardia nova* complex.[2] Authors explain that how identified *N. asteroides* of other species in *N. asteroides* complex?
  - ii. The authors have not mentioned of the use of molecular techniques to *Nocardia* identification at species level in the article. The authors explain molecular method if used.
3. In literature, although phenotypic methods are labor intensive and time-consuming, they are used in combination with molecular techniques such as polymerase chain reaction (PCR) sequencing (16S rRNA, *hsp65*, *rpoB*, *gyrB*, and *secA* genes) and PCR-restriction fragment length polymorphism for accurate identification in genus and species levels for *Nocardia*[2,13]
4. Drug choice for nocardiosis treatment is co-trimoxazole, but some of species are resistance to co-

trimoxazole and other antibacterial agents; therefore, accurate identification in species level and antimicrobial susceptibility testing are important.[2,14] Further, in literature, some of patients have hypersensitive reaction to co-trimoxazole.[15]

### Financial support and sponsorship

Nil.

### Conflicts of interest

There are no conflicts of interest.

### References

---

1. Shirani K, Poulsen AN, Hakamifard A. Nocardial brain abscess in a patient with pulmonary alveolar proteinosis. *Adv Biomed Res.* 2015;4:185. [PMCID: PMC4617009] [PubMed: 26605224]
2. Brown-Elliott BA, Brown JM, Conville PS, Wallace RJ, Jr Clinical and laboratory features of the *Nocardia* spp. based on current molecular taxonomy. *Clin Microbiol Rev.* 2006;19:259–82. [PMCID: PMC1471991] [PubMed: 16614249]
3. Goodfellow M. Characterisation of *Mycobacterium*, *Nocardia*, *Corynebacterium* and related taxa. *Ann Soc Belg Med Trop.* 1973;53:287–98. [PubMed: 4578844]
4. Goodfellow M, Lind A, Mordarska H, Pattyn S, Tsukamura M. A co-operative numerical analysis of cultures considered to belong to the ‘rhodochrous’ taxon. *Microbiology.* 1974;85:291–302.
5. Broughton RA, Wilson HD, Goodman NL, Hedrick JA. Septic arthritis and osteomyelitis caused by an organism of the genus *Rhodococcus*. *J Clin Microbiol.* 1981;13:209–13. [PMCID: PMC273751] [PubMed: 7462414]
6. Severo LC, Petrillo VF, Coutinho LM. Actinomycetoma caused by *Rhodococcus* spp. *Mycopathologia.* 1987;98:129–31. [PubMed: 3587337]
7. Arenskötter M, Bröker D, Steinbüchel A. Biology of the metabolically diverse genus *Gordonia*. *Appl Environ Microbiol.* 2004;70:3195–204. [PMCID: PMC427784] [PubMed: 15184112]
8. Blanc V, Dalle M, Markarian A, Debonne MV, Duplay E, Rodriguez-Nava V, et al. *Gordonia terrae*: A difficult-to-diagnose emerging pathogen? *J Clin Microbiol.* 2007;45:1076–7. [PMCID: PMC1829133] [PubMed: 17192419]
9. Blaschke AJ, Bender J, Byington CL, Korgenski K, Daly J, Petti CA, et al. *Gordonia* species: Emerging pathogens in pediatric patients that are identified by 16S ribosomal RNA gene sequencing. *Clin Infect Dis.* 2007;45:483–6. [PubMed: 17638199]
10. Prescott JF. *Rhodococcus equi*: An animal and human pathogen. *Clin Microbiol Rev.* 1991;4:20–34. [PMCID: PMC358176] [PubMed: 2004346]
11. Goodfellow M, Alderson G. The actinomycete-genus *Rhodococcus*: A home for the ‘rhodochrous’ complex. *Microbiology.* 1977;100:99–122.
12. Liu CY, Lai CC, Lee MR, Lee YC, Huang YT, Liao CH, et al. Clinical characteristics of infections caused by *Tsukamurella* spp. and antimicrobial susceptibilities of the isolates. *Int J Antimicrob Agents.* 2011;38:534–7. [PubMed: 22014886]
13. McTaggart LR, Richardson SE, Witkowska M, Zhang SX. Phylogeny and identification of *Nocardia* species on the basis of multilocus sequence analysis. *J Clin Microbiol.* 2010;48:4525–33. [PMCID: PMC3008441] [PubMed: 20844218]
14. Brown-Elliott BA, Biehle J, Conville PS, Cohen S, Saubolle M, Sussland D, et al. Sulfonamide resistance in isolates of *Nocardia* spp. from a US multicenter survey. *J Clin Microbiol.* 2012;50:670–2. [PMCID: PMC3295118] [PubMed: 22170936]

15. Walton N, Krumins D. Rapid desensitisation of an immunocompetent patient with cotrimoxazole hypersensitivity. *J Pharm Pract Res.* 2010;40:130–2.

---

Articles from Advanced Biomedical Research are provided here courtesy of **Medknow Publications**