



Vector-borne pathogens in dogs of different regions of Iran and Pakistan

Roberta Iatta¹ · Alireza Sazmand^{2,3} · Viet-Linh Nguyen¹ · Farzad Nemati² · Muhammad Mazhar Ayaz⁴ · Zahra Bahiraei² · Salman Zafari² · Anna Giannico⁵ · Grazia Greco¹ · Filipe Dantas-Torres⁶ · Domenico Otranto^{1,2} 

Received: 16 October 2020 / Accepted: 25 November 2020
© The Author(s) 2021

Abstract

Canine vector-borne diseases (CVBDs) are highly prevalent in tropical and subtropical countries, mainly due to favorable climate conditions and reduced adoption of preventive measures. This study aimed to provide a comprehensive overview on the prevalence of CVBDs in Iran and Pakistan where limited data are available. Blood samples were collected from 403 dogs from six provinces in Iran and Pakistan to assess the presence of pathogen DNA (i.e., *Anaplasma* spp., *Coxiella burnetii*, *Ehrlichia* spp., *Rickettsia* spp., *Babesia* spp., *Hepatozoon* spp., filarioids, and *Leishmania* spp.). Sera were also screened by an immunofluorescence antibody test for the detection of antibodies against *Leishmania infantum*. In total, 46.9% of dogs scored positive to *Hepatozoon canis* being the most frequently detected (41.4%), followed by *Anaplasma platys* (6.4%), *Ehrlichia canis* (3.4%), *Rickettsia* spp. (2.2%), *Babesia vogeli* (1.0%), and *L. infantum* (0.3%). A seroprevalence of 9.6% to anti-*L. infantum* IgG was also recorded. Data reported herein demonstrate that dogs from Iran and Pakistan are at a high risk of CVBDs, particularly of canine hepatozoonosis. Effective control strategies are advocated for minimizing the risk of infection in animals and humans, also in consideration of the zoonotic potential of some pathogens detected.

Keywords *Anaplasma platys* · Canine vector-borne pathogens · *Ehrlichia canis* · *Rickettsia* spp. · *Hepatozoon canis* · *Leishmania infantum* · Iran · Pakistan

Roberta Iatta and Alireza Sazmand contributed equally to this work.

Guest Editor: Christina Strube

✉ Domenico Otranto
domenico.otranto@uniba.it

- ¹ Department of Veterinary Medicine, University of Bari, Bari, Italy
- ² Department of Pathobiology, Faculty of Veterinary Science, Bu-Ali Sina University, Hamedan, Iran
- ³ Zoonotic Diseases Research Center, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
- ⁴ Department of Parasitology, Faculty of Veterinary Sciences, Cholistan University of Veterinary and Animal Sciences, Bahawalpur, Pakistan
- ⁵ Istituto Zooprofilattico della Puglia e della Basilicata, Putignano (BA), Italy
- ⁶ Department of Immunology, Aggeu Magalhães Institute, Oswaldo Cruz Foundation (Fiocruz), Recife, Brazil

Introduction

Vector-borne diseases (VBDs) are of growing concern to people across the world and their increasing incidence has been attributed to several factors, such as climate change and animal movements (Ogden and Lindsay 2016). Their distribution depends on a complex combination of biotic and abiotic factors, making their control extremely difficult (Dantas-Torres et al. 2012; Otranto et al. 2017). The impact of VBDs is heavier on tropical and subtropical countries, where the climate is more suitable for various arthropod vectors and where people and animals have limited access to healthcare services, including diagnosis and treatment of these diseases (Otranto et al. 2017; Colella et al. 2019).

Dogs serve as blood source to arthropod vectors and are suitable reservoirs of many vector-borne pathogens (VBPs; Otranto 2018). For example, dogs are well recognized as the primary reservoirs of *Leishmania infantum*, the causative agent of zoonotic visceral leishmaniasis and the target of integrated control strategies (Travi et al. 2018; Dantas-Torres et al. 2019). Also, dogs are primary reservoirs of some

mosquitoes-transmitted filarial helminths (e.g., *Dirofilaria immitis* and *Dirofilaria repens*), which may pose a risk to humans in areas where dogs are highly infected, including municipal shelters (Panarese et al. 2020). In the latter case, dog relocation projects may also contribute to increase pathogen circulation in previously non-endemic regions, and eventually their transmission, when a proper vector is present (Otranto et al. 2017; Mendoza-Roldan et al. 2020).

Dogs also contribute to the circulation of certain tick species, such as *Rhipicephalus sanguineus* sensu lato (s.l.), as well as other more generalist tick species (e.g., *Ixodes ricinus*) which are well recognized as vectors of pathogens to animals and humans (Otranto et al. 2009a).

Therefore, canine vector-borne diseases (CVBDs) caused by a wide range of pathogens, including viruses, bacteria, protozoa, and helminths, are of veterinary importance and may represent a public health issue (Otranto et al. 2009a, 2017). Data on the occurrence of CVBDs are reported in a few countries of the Middle East such as Iraq (Otranto et al. 2019), Turkey (Kirkan et al. 2013), and Qatar (Alho et al. 2017), making their impact on animal and human populations difficult to estimate, which ultimately impairs the implementation of preventive strategies for minimizing the risk of infections. For example, the occurrence of VBPs has been documented only in a few regions and on limited number of dogs in Iran and Pakistan (Khazeni et al. 2013; Khedri et al. 2014; Motaghpisheh et al. 2016; Ahmad et al. 2018a; Mohebbali et al. 2018; Barati and Razmi 2018; Azari-Hamidian et al. 2019). In this perspective, the aim of this study was to investigate the prevalence of VBPs in sheltered and owned dogs from five Iranian provinces and from Punjab in Pakistan in order to fill gaps in the knowledge of VBPs in these areas.

Materials and methods

Sample collection

From October 2018 to November 2019, blood samples (1.5–5 ml) were collected from cephalic or saphenous vein of 403 dogs (i.e., 357 sheltered and 46 privately owned dogs) from 5 cities in 5 provinces of Iran including Amol in the North ($n = 75$), Hamedan ($n = 81$) and Kermanshah ($n = 51$) in the west, Yazd in the center ($n = 78$), Ahvaz in the south-west ($n = 69$), and from Bahawalpur in the eastern province of Punjab, Pakistan ($n = 49$) (Fig. 1). Animal data (i.e., age, sex, breed, neutering status, and province) were recorded. All dogs stayed outside overnight. Dogs were grouped according to age in ≤ 1 year (G1), >1 to <5 years (G2), and ≥ 5 years (G3). The blood of dogs was collected with permission of the Ethical Committee of Hamedan University of Medical Sciences, Iran (code: IR.UMSHA.REC.1398.124).

Serological testing

Serum samples from 354 out of 403 dogs were tested for anti-*L. infantum* antibodies by using an immunofluorescence antibody test (IFAT), as described elsewhere (Otranto et al. 2009c). Promastigotes of *L. infantum* zymodeme MON-1 were used as antigen. Samples were scored as positive when they produced a clear cytoplasmic and membrane fluorescence of promastigotes at a cut-off dilution of 1:80. Positive sera were titrated by serial dilutions until negative results were obtained.

DNA extraction, PCR protocols, and sequencing

Genomic DNA was extracted from 200 μ l aliquots of EDTA-treated blood samples using GenUP DNA Kit (Biotechrabbit, Berlin, Germany) following the manufacturer's instructions. All samples were tested for the presence of *Anaplasma* spp., *Ehrlichia* spp., *Rickettsia* spp., *Babesia* spp., *Hepatozoon* spp., and filarioid DNA by using conventional PCR (cPCR). In particular, for the detection of *Rickettsia* spp., DNA samples were firstly tested by PCR targeting a partial sequence of the gene citrate synthase (*gltA*) and positive samples were further tested by a second PCR targeting a fragment of the outer membrane protein A (*ompA*) gene. All primers and cPCR protocols used for the detection of VBPs are summarized in Table 1. *Leishmania* spp. and *Coxiella burnetii* were detected by using real-time PCR (qPCR) and positive samples were further tested by cPCR. For all reactions, DNA of pathogen-positive blood samples served as a positive control. Amplified PCR products were visualized in 2% agarose gel stained with GelRed (VWR International PBI, Milan, Italy) by GelLogic 100 gel documentation system (Kodak, NY, USA). The cPCR amplicons were sequenced using the Big Dye Terminator v.3.1 chemistry in a 3130 Genetic Analyzer (Applied Biosystems, CA, USA). Nucleotide sequences were edited, aligned, and analyzed using the Geneious platform version 7.0 (Biomatters Ltd., Auckland, New Zealand) (Kearse et al. 2012) and compared with those available in the GenBank® database using the Basic Local Alignment Search Tool (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Phylogenetic analysis

For phylogenetic analyses, the 18S rRNA and *gltA* representative sequence types (STs) of *Hepatozoon* spp. and *Rickettsia* spp., respectively, and the corresponding sequences available from the GenBank® database were included. Phylogenetic relationships were inferred using the maximum likelihood (ML) method based on Hasegawa-Kishino-Yano model (Hasegawa et al. 1985) for *Hepatozoon canis* analysis and Tamura 3-parameter model (Tamura 1992) for *Rickettsia* spp. Gamma distribution (+G) was used to model evolutionary rate differences among sites selected by best-fit model




Fig. 1 Samples were collected from different provinces in Iran and Pakistan

(Nei and Kumar 2000). Evolutionary analyses were conducted on 8000 bootstrap replications using the MEGA X software (Kumar et al. 2018). Homologous sequences from *Adelina bambarooniae* (accession nos. AF494058) as well as *Rickettsia typhi* and *Rickettsia prowazekii* (accession nos. U59714 and U59715) were used as outgroups for *Hepatozoon* spp. and *Rickettsia* spp. phylogenetic trees, respectively.

Statistical analysis

Exact binomial 95% confidence intervals (CIs) were established for proportions. The chi-square and Fisher tests were used to compare proportions with a probability p value < 0.05 regarded as statistically significant. Analyses were done using the GraphPad Prism version 8.0.0 (GraphPad Software, San Diego, CA, USA).

Results

Out of 403 dogs tested, 189 (46.9%; 95% CI 42.1–51.9) scored positive to at least one pathogen. Data on sex, age, and location of animals are reported in Table 2 along with number and percentage of dogs positive to *H. canis*, the most frequently detected pathogen species (41.4%; 95% CI 36.7–46.3) and seropositive to antibodies against *L. infantum* (9.6%; 95% CI 6.9–13.1). The prevalence of infection by other pathogens including *Anaplasma platys*, *Ehrlichia canis*,

spotted-fever group (SFG) *Rickettsia* spp., *Babesia* spp., and *L. infantum* is reported in Table 3. No DNA of filarioids or *C. burnetii* was amplified. Co-infections with *H. canis* and *A. platys* were the most frequent (4.9%, $n = 20$), followed by *H. canis* and *E. canis* (1.4%, $n = 3$); *H. canis* and *Rickettsia* spp. (1.4%, $n = 3$); *H. canis* and *L. infantum* (0.3%, $n = 1$); and *H. canis*, *E. canis*, and *Rickettsia* spp. (0.3%, $n = 1$). The *ompA* gene was not amplified in any of the samples positive to *Rickettsia gltA* gene. Out of 354 serum samples, 34 (9.6%) were positive to anti-*L. infantum* antibodies with titers of 1:80 ($n = 14$), 1:160 ($n = 15$), 1:320 ($n = 3$), 1:640 ($n = 1$), and 1:1280 ($n = 1$).

The risk of *H. canis* infection in dogs was significantly associated with their keeping conditions and the geographical areas, whereas the risk of *L. infantum* exposure only with the geographical areas. In particular, sheltered dogs were related to higher risk of *H. canis* infection than owned dogs ($\chi^2 = 29.4$, $df = 1$, $p < 0.00001$). Based on location, the highest prevalence of *H. canis* infection (i.e., 78.4%) was recorded in Kermanshah in Iran followed by Bahawalpur in Pakistan (63.3%) and both percentages were significantly higher than those recorded in other cities ($p < 0.05$). Dogs from Amol in Iran were more exposed to *L. infantum* than those in the other Iranian provinces ($p < 0.05$) (Table 2). No statistical association was found between *H. canis* infection or *L. infantum* exposure in dogs and their sex and age. The region with the highest pathogen diversity was Ahvaz in Iran where a high prevalence of *Rickettsia* spp. infection was recorded (13%) (Table 3).

Table 1 Primers, target genes, and PCR conditions used in this study

Pathogen	Primers	Target gene	Product size (bp)	Reference
<i>Ehrlichia</i> spp./ <i>Anaplasma</i> spp.	EHR16SD: GGTACCYACAGAAGAAGTCC EHR16SR: TAGCACTCATCGTTTACA GC	16S rRNA	345	Martin et al. (2005)
<i>Babesia</i> spp./ <i>Hepatozoon</i> spp.	Piroplasmid-F: CCAGCAGCCGCGTAAATTC Piroplasmid-R: CTTTCGCAGTAGTTYGTCTTTAACAA ATCT	18S rRNA	350–400	Tabar et al. (2008)
<i>Rickettsia</i> spp.	CS-78F: GCAAGTATCGGTGAGGATGTAAT CS-323R: GCTTCCTTAAAAATTCAATAAATCAGGAT Rr190.70F: ATGGCGAATATTTCTCCAAAA Rr190.701R: GTTCCGTTAATGGCAGCATCT	<i>gltA</i> <i>opmA</i>	401 632	Labruna et al. (2004) Regnery et al. (1991)
<i>Coxiella burnetii</i>	Cox-F: GTCTTAAGGTGGGCTGCGTG Cox-R: CCCCGAATCTCATTGATCAGC Cox-TM: FAM-AGCGAACCATTGGTATCGGA CGTT-TAMRA-TATGG	<i>IS1111</i>	295	Klee et al. (2006)
<i>Leishmania</i> spp.	LGITSF2: GCATGCCATATTCTCAGTGTC LGITSR2: GGCCAACGCGAAGTTGAATTC LEISH-1: AACTTTTCTGGTCCCTCCGGGTAG LEISH-2: ACCCCCAGTTTCCCGCC Probe: FAM-AAAAATGGGTGCAGAAAT	ITS-2 kDNA minicircle	383–450 120	de Almeida et al. (2011) Francino et al. (2006)
Nematodes	NTF: TGATTGGTGGTTTTGGTAA NTR: ATAAGTACGAGTATCAATATC	<i>cox1</i>	648	Casiraghi et al. (2001)

Consensus sequences for each detected pathogen displayed 99–100% nucleotide identity with those available in the GenBank® database. In particular, *Babesia vogeli* ($n = 4$) nucleotide similarity was 99.3–99.6% with MT499357. In addition, five STs were identified for *H. canis* (ST1, $n = 6$,

100% identity with MN393911; ST2, $n = 10$, 99.7% with MK673827; ST3, $n = 24$, 100% with KX880505; ST4, $n = 39$, 100% with MT499354; ST5, $n = 88$, 100% with MK673850) and *L. infantum* 100% identity was recorded with MK496879. *Anaplasma platys* sequences ($n = 26$) and

Table 2 Number and percentage of dogs positive to *Hepatozoon canis* DNA ($n = 403$) and antibodies against *Leishmania infantum* ($n = 354$) according to their sex, age, keeping condition, and sampling area. (NA indicates data not available)

Variables	<i>Hepatozoon canis</i> Pos/total (%; 95% CI)	<i>Leishmania infantum</i> Pos/total (%; 95% CI)
Sex		
Male	73/161 (45.3; 37.6–53.1)	12/122 (9.8; 5.6–16.7)
Female	92/240 (38.3; 32.3–44.8)	22/230 (9.6; 6.2–14.1)
No data	2/2 (100)	0/2 (0)
Age		
≤ 1 year	34/84 (40.5; 30.3–51.2)	4/59 (6.8; 2.3–16.7)
> 1 year to < 5 years	96/241 (39.8; 33.8–46.3)	18/219 (8.2; 5.2–12.7)
≥ 5 years	34/74 (45.9; 34.6–57.5)	12/74 (16.2; 9.3–26.3)
No data	3/4 (75.0; 24.8–98.2)	0/2 (0)
Keeping condition		
Privately owned	2/46 (4.3; 0.8–14.9)	8/46 (17.4; 9.0–30.7)
Shelter	165/357 (46.2; 41–51.5)	26/308 (8.4; 5.7–12.1)
Geographical origin (country/city)		
Iran		
Ahvaz	34/69 (49.3; 37.6–60.9)	6/69 (8.7; 3.8–17.9)
Amol	32/75 (42.7; 31.9–54.1)	19/75 (25.3; 16.5–36.6)
Kermanshah	40/51 (78.4; 64.8–87.9)	0/51 (0)
Hamedan	15/81 (18.5; 11.2–28.3)	9/81 (11.1; 5.7–20.2)
Yazd	15/78 (19.2; 11.7–29.4)	0/78 (0)
Pakistan		
Bahawalpur	31/49 (63.3; 48.9–75.7)	NA
Total	167/403 (41.4; 36.7–46.4)	34/354 (9.6; 6.9–13.1)

Table 3 Number (percentage) of dogs molecularly positive to vector-borne pathogens according to their location

Country	Province	Total number of dogs infected with each pathogen (%)				
		<i>Anaplasma platys</i>	<i>Ehrlichia canis</i>	<i>Rickettsia</i> spp.	<i>Babesia vogeli</i>	<i>Leishmania infantum</i>
Iran	Ahvaz (69)	1 (1.4)	1 (1.4)	8 (13) [^]		
	Amol (75)	2 (2.7)	1 (1.3)			
	Hamedan (81)	-			4 (4.9)	
	Kermanshah (51)	10 (19.6)				1 (2)
	Yazd (78)	-	-			
Pakistan	Bahawalpur (49)	13 (26.5)	12 (24.5)	1 (2) [°]		
Total (403)		26 (6.4)	14 (3.4)	9 (2.2)	4 (0.99)	1 (0.25)

[^]*n* = 4, *Rickettsia monacensis*; *n* = 2, *Rickettsia helvetica*; *n* = 2, *Rickettsia heilongjiangensis*/*Rickettsia raoultii*/*Rickettsia slovacica*

[°]*n* = 1, *Rickettsia conorii*/*Rickettsia honei*/*Rickettsia raoultii*

E. canis (*n* = 14) had 100% nucleotide identity with MN922611 and MN922610, respectively. As far as *Rickettsia* spp., four samples had 100% nucleotide identity with *Rickettsia monacensis* (KC993860) and two had 99.7% identity with *Rickettsia helvetica* (U59723). One sample from a dog in Pakistan was 100% identical to *Rickettsia conorii* (U59730)/*Rickettsia honei* (U59726)/*Rickettsia raoultii* (CP019435), and two other samples from dogs in Ahvaz (Iran) were 100% identical to *Rickettsia heilongjiangensis* (CP002912)/*Rickettsia raoultii* (DQ365803)/*Rickettsia slovacica* (U59725).

Molecular identification of nucleotide sequences for *Rickettsia* spp. and *H. canis* was supported by the distinct separation of species-specific clades inferred from the phylogenetic analyses. The ML tree of *H. canis* grouped all representative STs in a large clade supported by high bootstrap value (i.e., 98%), to the exclusion of other species of *Hepatozoon* (Fig. 2). The ML tree based on the partial *gltA* gene sequences of *Rickettsia* spp. showed that all sequences, herein detected, clustered in well-supported clades with other SFG rickettsiae (Fig. 3). Representative sequences of pathogens detected in this study were deposited in the GenBank® database under the accession numbers MW019628 and MW019629 for *B. vogeli*, MW019630-MW019643 for *H. canis*, MW019670 and MW019671 for *A. platys*, MW019672 and MW019673 for *E. canis*, MW039480 for *R. helvetica*, MW039481 for *R. monacensis*, MW039482 and MW039483 for *Rickettsia* spp., and MW074300 for *L. infantum*.

Discussion

The high frequency of CVBD-causing pathogens reported herein indicates that dog populations from Iran and Pakistan are exposed to multiple pathogens, including some of

zoonotic importance, thus posing a risk not only to dogs, but also to public health. Many of the herein detected pathogens (i.e., *A. platys*, *E. canis*, *B. vogeli*, and *H. canis*) are transmitted by *Rh. sanguineus* s.l. (Latrofa et al. 2014), the most common tick species found on dogs in Iran and Pakistan (Mosallanejad et al. 2012; Cabezas-Cruz et al. 2019). The high prevalence of *H. canis* in all the provinces of Iran as well as from Bahawalpur in Pakistan suggests that *Rh. sanguineus* s.l. is highly prevalent in these areas. After the first detection of *H. canis* gametocytes in a blood smear of an Iranian dog (Khoshnegah et al. 2009), the prevalence of this infection was reported only in dogs from northern half of Iran ranging from 1.6 to 23% (Amoli et al. 2012; Dalimi et al. 2017; Soltani and Dalimi 2018; Barati and Razmi 2018). Conversely, the prevalence herein recorded for *H. canis* in Pakistan (i.e., 41.4%) was similar to that of farm dogs recorded in Punjab districts (45.5%, Ahmad et al. 2018a). Scientific data indicates that under certain circumstances, such as in poor socioeconomic settings, the high environmental infestation of arthropod vectors and the absence of preventive strategies in dogs and in the environment concur to increase the incidence of CVBD (Otranto et al. 2017). Indeed, similar high prevalence of *H. canis* infections has been reported in dogs (37.8%) and in *Rh. sanguineus* s.l. ticks (42.5%) from India (Manoj et al. 2020) as well as in dogs from Iraq (33%) (Otranto et al. 2019). Although *Rh. sanguineus* s.l., *Rhipicephalus bursa* and *Rhipicephalus annulatus* are the common species infesting dogs in the studied area (Jamshidi et al. 2012; Akhtardanesh et al. 2016; Ali et al. 2019; Cabezas-Cruz et al. 2019), the circulation of other competent tick vectors cannot be ruled out (Murata et al. 1995; Rubini et al. 2009; Najm et al. 2014; Giannelli et al. 2017). In addition, the significantly higher occurrence of dogs infected by *H. canis* in Kermanshah in Iran (78.4%) and Bahawalpur in Pakistan (63.3%) could be related to the high population density of tick vectors in these shelters observed during the dog sampling.




Fig. 2 Phylogenetic relationship of *Hepatozoon* spp. sequences isolated in this study (in bold) to other *Hepatozoon* spp. based on a partial sequence (327 bp) of the 18S rRNA gene. The analyses were performed using a maximum likelihood with Hasegawa-Kishino-Yano

model. A gamma distribution was used to model evolutionary rate differences among sites. Homologous sequence from *Adelina bambarooniae* (accession nos. AF494058) was used as the outgroup

The risk for sheltered dogs to be more infected by *H. canis* than owned dogs suggests that in this environment, the animals are more exposed to the vectors. Indeed, preventive measures such as ectoparasite repellents are not commonly

employed in sheltered dogs in Iran and Pakistan because of the limitation in budget of non-governmental organization which mainly aim to feed and protect dogs from culling program (personal observations of AS and MMA). Conversely,

Fig. 3 Phylogenetic relationship of *Rickettsia* spp. sequences isolated in this study (in bold) to other *Rickettsia* strains based on a partial sequence (345 bp) of the *gltA* gene. The analyses were performed using a maximum likelihood method with Tamura 3-parameter model. A gamma distribution was used to model evolutionary rate differences among sites. Homologous sequences from *Rickettsia typhi* (accession nos. U59714) and *Rickettsia prowazekii* (accession nos. U59715) were used as the outgroups



B. vogeli was found in only few dog samples (i.e., 1%) as recorded in other studies from Iran (Alborzi et al. 2013; Akhtardanesh et al. 2016; Bigdeli and Namavari 2017) and Pakistan (Ahmad et al. 2018b). Also, *Babesia gibsoni* was

reported causing canine babesiosis in these two countries (Akhtardanesh et al. 2016; Ahmad et al. 2018b).

In Iran, the first study on leishmaniosis in carnivores was conducted in 1982 with a seroprevalence of 2.4% in jackals

and 3% in dogs from the northern part (Hamidi et al. 1982). Afterwards, *L. infantum* infection was reported in domestic and wild animals in most of the regions of Iran with differences in frequency probably related to geographic regions, environmental conditions, and dog population (Shokri et al. 2017; Mohebbali et al. 2018). The seropositivity to *L. infantum* in dogs from Iran herein detected (9.6%) confirms the circulation of infected sand fly vector (Yaghoobi-Ershadi 2016), thus the risk of dogs as well as humans to be exposed to their bites and to be infected. Differences in climatic factors may affect the sand fly population (Cheghabalaki et al. 2019) in terms of species composition and abundance, thus leading to a higher risk of infection in dogs living in some provinces. Indeed, the highest seroprevalence (25.3%) was recorded in dogs from Amol, located in the north of Iran with a Mediterranean climate vs no infected dog in Yazd and Kermanshah which are regions characterized by hot and dry climate. The detection of *L. infantum* DNA only in one dog may also be related to the use of the blood, which is a convenient type of sample but poor as a parasite source (Maia et al. 2009; Otranto et al. 2009c).

Canine monocytic ehrlichiosis by *E. canis* and canine cyclic thrombocytopenia by *A. platys* affect dogs worldwide, varying from asymptomatic infections or with mild symptoms to a severe illness (Otranto et al. 2009b). Although the prevalence of these infections is scantily reported in the examined areas, the finding of *E. canis* in dogs from Iran and Pakistan provinces is not surprising since this bacterium is transmitted by *Rh. sanguineus* s.l. A few seroprevalence studies (Akhtardanesh et al. 2010) as well as molecular detection of *E. canis* (Motaghipisheh et al. 2016; Malik et al. 2018) in blood samples confirm that dogs from the examined areas are exposed to the pathogen. Conversely, the first report of *A. platys* in both countries with higher prevalence than *E. canis* (6.4% vs 3.4%) indicates that this pathogen may affect dogs living mainly in rural area and in shelters. While the unsuccessful amplification of the *ompA* gene may represent a limitation of the study, some sequences obtained from dogs from Iran had high (> 99%) sequence identity with *R. helvetica* and *R. monacensis*. Although the pathogenicity of *R. helvetica* and *R. monacensis* in dogs is unknown, both species may cause severe diseases in humans as reported in Europe and Southeast Asia for *R. helvetica* (Nilsson et al. 1999; Fournier et al. 2004; Nilsson et al. 2010) and in Spain and Italy for *R. monacensis* (Jado et al. 2007; Madeddu et al. 2012). Data on SFG rickettsiae in dogs and their ectoparasites in the Middle East are scant (Baneth et al. 1998; Harrus et al. 2011; Levin et al. 2011; Kirkan et al. 2013; Parola et al. 2013; Orkun et al. 2014). Our findings suggest the occurrence of SFG rickettsiae in dogs from Iran and Pakistan, but further studies are needed to confirm the identity of these microorganisms and also to understand the potential risks for public health.

In this study, none of the tested dogs scored positive to *C. burnetii* although this bacteria was serologically diagnosed in a domestic dog in Ahvaz, Iran (Rezaei et al. 2016). Indeed, Q fever is endemic in Iran and Pakistan with high prevalence among human patients and domestic ruminants (Ullah et al. 2019; Esmaeili et al. 2019; Mohabbati Mobarez et al. 2017). In fact, it is acknowledged that ruminants (cattle, sheep, and goats) are the most important reservoirs of *C. burnetii* for human infection (Angelakis and Raoult 2010).

In conclusion, the herein reported data provide more knowledge of CVBDs in these countries suggesting that different pathogens as well as arthropod vectors circulate among animal populations. Of the detected pathogens, SFG rickettsiae are of great relevance for their pathogenicity to humans. Considering the occurrence of infection by zoonotic pathogens in dogs and their close relationship with humans, effective control strategies are advocated for minimizing the risk of infection in animals as well as in humans.

Acknowledgments This study was planned under the academic agreement between the Bu-Ali Sina University Hamedan (Iran) and the University of Bari (Italy). We thank all who assisted us in sampling especially Dr. Seyedmasoud Zolhavarieh (Bu-Ali Sina University), Dr. Omid Hosseinejad and Dr. Fatemeh Arabifar (in Ahvaz), Mrs. Shilan Lorestani and Dr. Saman Salmani (in Kermanshah), Mrs. Faranak Rahmanpour (in Amol), and Mr. Seyed-Ali Vaziri (in Yazd).

Funding Open Access funding provided by Università degli Studi di Bari Aldo Moro. This work was partially supported by the Bu-Ali Sina University, Hamedan, Iran (Grant Number 1/1/28864).

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ahmad AS, Saeed MA, Rashid I et al (2018a) Molecular characterization of *Hepatozoon canis* from farm dogs in Pakistan. Parasitol Res 117: 1131–1138. <https://doi.org/10.1007/s00436-018-5790-1>
- Ahmad AS, Rashid I, Ashraf K et al (2018b) Molecular occurrence of canine babesiosis in rural dog population in Pakistan. Trop Biomed 35:593–603

- Akhtardanesh B, Ghanbarpour R, Blourizadeh H (2010) Serological evidence of canine monocytic ehrlichiosis in Iran. *Comp Clin Pathol* 19:469–474. <https://doi.org/10.1007/s00580-009-0889-5>
- Akhtardanesh B, Saberi M, Nurollahifard SR, Aghazamani M (2016) Molecular detection of *Babesia* spp. in tick-infested dogs in south-eastern Iran. *J Dis Glob Heal* 8:72–77
- Alborzi AR, Avizeh R, Mosallanejad B et al (2013) *Babesia* infection in urban and rural dogs in Ahvaz district, Southwest of Iran. *Arch Razi Inst* 68:37–42. <https://doi.org/10.7508/ari.2013.01.006>
- Alho AM, Lima C, Latrofa MS et al (2017) Molecular detection of vector-borne pathogens in dogs and cats from Qatar. *Parasit Vectors* 10:1–5. <https://doi.org/10.1186/s13071-017-2237-y>
- Ali A, Khan MA, Zahid H et al (2019) Seasonal dynamics, record of ticks infesting humans, wild and domestic animals and molecular phylogeny of *Rhipicephalus microplus* in Khyber Pakhtunkhwa Pakistan. *Front Physiol* 10:1–15. <https://doi.org/10.3389/fphys.2019.00793>
- Amoli AR, Khoshnegah J, Razmi G (2012) A preliminary parasitological survey of *Hepatozoon* spp. infection in dogs in Mashhad, Iran. *Iran J Parasitol* 7:99–103
- Angelakis E, Raoult D (2010) Q fever. *Vet Microbiol* 140:297–309. <https://doi.org/10.1016/j.vetmic.2009.07.016>
- Azari-Hamidian S, Norouzi B, Harbach RE (2019) A detailed review of the mosquitoes (Diptera: Culicidae) of Iran and their medical and veterinary importance. *Acta Trop* 194:106–122. <https://doi.org/10.1016/j.actatropica.2019.03.019>
- Baneth G, Breitschwerdt EB, Hegarty BC et al (1998) A survey of tick-borne bacteria and protozoa in naturally exposed dogs from Israel. *Vet Parasitol* 74:133–142. [https://doi.org/10.1016/S0304-4017\(97\)00149-0](https://doi.org/10.1016/S0304-4017(97)00149-0)
- Barati A, Razmi GR (2018) A parasitologic and molecular survey of *Hepatozoon canis* infection in stray dogs in Northeastern Iran. *J Parasitol* 104:413–417. <https://doi.org/10.1645/17-105>
- Bigdeli M, Namavari M (2017) Evaluation of PCR assay using specific primers in diagnosis of canine ehrlichiosis and babesiosis: a study on herd and stray dogs in Shiraz. *J Altern Vet Med* 1:1–15
- Cabezas-Cruz A, Allain E, Ahmad AS et al (2019) Low genetic diversity of *Ehrlichia canis* associated with high co-infection rates in *Rhipicephalus sanguineus* (s.l.). *Parasit Vectors* 12:1–13. <https://doi.org/10.1186/s13071-018-3194-9>
- Casiraghi M, Anderson TJC, Bandi C et al (2001) A phylogenetic analysis of filarial nematodes: comparison with the phylogeny of *Wolbachia* endosymbionts. *Parasitology* 122:93–103. <https://doi.org/10.1017/S0031182000007149>
- Cheghabalaki ZZ, Yarahmadi D, Karampour M, Shamsipour A (2019) Spatial dynamics of a phlebotomine sand flies population in response to climatic conditions in Bushehr Province of Iran. *Ann Glob Heal* 85:1–11. <https://doi.org/10.5334/aogh.30>
- Colella V, Hodžić A, Iatta R et al (2019) Zoonotic Leishmaniasis, Bosnia and Herzegovina. *Emerg Infect Dis* 25:385–386. <https://doi.org/10.3201/eid2502.181481>
- Dalimi A, Jameie F, Mohammadi A et al (2017) Molecular detection of *Hepatozoon canis* in dogs of Ardabil Province, Northwest of Iran. *Arch Razi Inst* 72:197–201. <https://doi.org/10.22034/ari.2017.108389>
- Dantas-Torres F, Chomel BB, Otranto D (2012) Ticks and tick-borne diseases: a one health perspective. *Trends Parasitol* 28:437–446. <https://doi.org/10.1016/j.pt.2012.07.003>
- Dantas-Torres F, Miró G, Baneth G et al (2019) Canine leishmaniasis control in the context of one health. *Emerg Infect Dis* 25:E1–E4. <https://doi.org/10.3201/eid2512.190164>
- de Almeida M, Steurer F, Koru O et al (2011) Identification of *Leishmania* spp. by molecular amplification and DNA sequencing analysis of a fragment of rRNA internal transcribed spacer 2. *J Clin Microbiol* 49:3143–3149. <https://doi.org/10.1128/JCM.01177-11>
- Esmaeili S, Bagheri Amiri F, Mokhayeri H et al (2019) Seroepidemiological study of Q fever, brucellosis and tularemia in butchers and slaughterhouses workers in Lorestan, western of Iran. *Comp Immunol Microbiol Infect Dis* 66:101322. <https://doi.org/10.1016/j.cimid.2019.06.003>
- Fournier P-E, Allombert C, Supputamongkol Y et al (2004) Aneruptive fever associated with antibodies to *Rickettsia helvetica* in Europe and Thailand. *J Clin Microbiol* 42:816–818. <https://doi.org/10.1128/jcm.42.2.816-818.2004>
- Francino O, Altet L, Sánchez-Robert E et al (2006) Advantages of real-time PCR assay for diagnosis and monitoring of canine leishmaniasis. *Vet Parasitol* 137:214–221. <https://doi.org/10.1016/j.vetpar.2006.01.011>
- Giannelli A, Lia RP, Annoscia G et al (2017) *Rhipicephalus turanicus*, a new vector of *Hepatozoon canis*. *Parasitology* 144:730–737. <https://doi.org/10.1017/S003118201600250X>
- Hamidi AN, Nadim A, Edrissian GH et al (1982) Visceral leishmaniasis of jackals and dogs in northern Iran. *Trans R Soc Trop Med Hyg* 76:756–757. [https://doi.org/10.1016/0035-9203\(82\)90100-6](https://doi.org/10.1016/0035-9203(82)90100-6)
- Harrus S, Perlman-Avrahami A, Mumcuoglu KY et al (2011) Molecular detection of *Rickettsia massiliae*, *Rickettsia sibirica mongolitimonae* and *Rickettsia conorii israelensis* in ticks from Israel. *Clin Microbiol Infect* 17:176–180. <https://doi.org/10.1111/j.1469-0691.2010.03224.x>
- Hasegawa M, Kishino H, Yano T (1985) Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *J Mol Evol* 22:160–174. <https://doi.org/10.1007/BF02101694>
- Jado I, Oteo JA, Aldámiz M et al (2007) *Rickettsia monacensis* and human disease, Spain. *Emerg Infect Dis* 13:1405–1407. <https://doi.org/10.3201/eid1309.060186>
- Jamshidi S, Maazi N, Ranjbar-Bahadori S et al (2012) A survey of ectoparasite infestation in dogs in Tehran, Iran. *Rev Bras Parasitol Vet* 21:326–329. <https://doi.org/10.1590/s1984-29612012000300030>
- Kearse M, Moir R, Wilson A et al (2012) Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>
- Khazeni A, Telmadarraiy Z, Oshaghi MA et al (2013) Molecular detection of *Ehrlichia canis* in ticks population collected on dogs in Meshkin-Shahr, Ardebil Province, Iran. *J Biomed Sci Eng* 6:1–5
- Khedri J, Radfar MH, Borji H et al (2014) Canine heartworm in Southeastern of Iran with review of disease distribution. *Iran J Parasitol* 9:560–567
- Khoshnegah J, Mohri M, Movassaghi AR, Mehrjerdi HK (2009) The first report of *Hepatozoon canis* infection of a dog in Iran. *Comp Clin Pathol* 18:455–458. <https://doi.org/10.1007/s00580-008-0794-3>
- Kirkan Ş, Savaşan S, Erbaş G, Parin U (2013) Prevalence of *Rickettsia rickettsii* infection in dogs from the urban and rural areas of western Turkey. *Ankara Üniv Vet Fak Derg* 60:165–169
- Klee SR, Tyczka J, Ellerbrok H et al (2006) Highly sensitive real-time PCR for specific detection and quantification of *Coxiella burnetii*. *BMC Microbiol* 6:2. <https://doi.org/10.1186/1471-2180-6-2>
- Kumar S, Stecher G, Li M et al (2018) MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* 35:1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Labruna MB, Whitworth T, Horta MC et al (2004) *Rickettsia* species infecting *Amblyomma cooperi* ticks from an area in the state of São Paulo, Brazil, where Brazilian spotted fever is endemic. *J Clin Microbiol* 42:90–98. <https://doi.org/10.1128/jcm.42.1.90-98.2004>
- Latrofa MS, Dantas-Torres F, Giannelli A, Otranto D (2014) Molecular detection of tick-borne pathogens in *Rhipicephalus sanguineus* group ticks. *Ticks Tick Borne Dis* 5:943–946. <https://doi.org/10.1016/j.ttbdis.2014.07.014>
- Levin ML, Killmaster LF, Zemtsova GE (2011) Domestic dogs (*Canis familiaris*) as reservoir hosts for *Rickettsia conorii*. *Vector-Borne Zoonotic Dis* 12:28–33. <https://doi.org/10.1089/vbz.2011.0684>

- Madeddu G, Mancini F, Caddeo A et al (2012) *Rickettsia monacensis* as cause of Mediterranean spotted fever-like illness, Italy. *Emerg Infect Dis* 18:702–704. <https://doi.org/10.3201/eid1804.111583>
- Maia C, Ramada J, Cristóvão JM et al (2009) Diagnosis of canine leishmaniasis: conventional and molecular techniques using different tissues. *Vet J* 179:142–144. <https://doi.org/10.1016/j.tvjl.2007.08.009>
- Malik MI, Qamar M, Ain Q et al (2018) Molecular detection of *Ehrlichia canis* in dogs from three districts in Punjab (Pakistan). *Vet Med Sci* 4:126–132. <https://doi.org/10.1002/vms3.94>
- Manoj RRS, Iatta R, Latrofa MS et al (2020) Canine vector-borne pathogens from dogs and ticks from Tamil Nadu, India. *Acta Trop* 203: 105308. <https://doi.org/10.1016/j.actatropica.2019.105308>
- Martin AR, Brown GK, Dunstan RH, Roberts TK (2005) *Anaplasma platys*: an improved PCR for its detection in dogs. *Exp Parasitol* 109:176–180. <https://doi.org/10.1016/j.exppara.2004.11.007>
- Mendoza-Roldan J, Benelli G, Panarese R et al (2020) *Leishmania infantum* and *Dirofilaria immitis* infections in Italy, 2009 – 2019: changing distribution patterns. *Parasit Vectors* 13:193. <https://doi.org/10.1186/s13071-020-04063-9>
- Mohabbati Mobarez A, Bagheri Amiri F, Esmaeili S (2017) Seroprevalence of Q fever among human and animal in Iran: a systematic review and meta-analysis. *PLoS Negl Trop Dis* 11: e0005521. <https://doi.org/10.1371/journal.pntd.0005521>
- Mohebbali M, Moradi-Asl E, Rassi Y (2018) Geographic distribution and spatial analysis of *Leishmania infantum* infection in domestic and wild animal reservoir hosts of zoonotic visceral leishmaniasis in Iran: a systematic review. *J Vector Borne Dis* 55:184–188. <https://doi.org/10.4103/0972-9062.249125>
- Mosallanejad B, Alborzi A, Katvandi N (2012) A survey on ectoparasite infestations in companion dogs of Ahvaz district, south-west of Iran. *J Arthropod Borne Dis* 6:70–78
- Motaghipisheh S, Akhtardanesh B, Ghanbarpour R et al (2016) Ehrlichiosis in household dogs and parasitized ticks in Kerman-Iran: preliminary zoonotic risk assessment. *J Arthropod Borne Dis* 10:246–252
- Murata T, Inoue M, Taura Y et al (1995) Detection of *Hepatozoon canis* oocyst from ticks collected from the infected dogs. *J Vet Med Sci* 57:111–112
- Najm N-A, Meyer-Kayser E, Hoffmann L et al (2014) *Hepatozoon canis* in German red foxes (*Vulpes vulpes*) and their ticks: molecular characterization and the phylogenetic relationship to other *Hepatozoon* spp. *Parasitol Res* 113:2679–2685. <https://doi.org/10.1007/s00436-014-3923-8>
- Nei M, Kumar S (2000) Molecular evolution and phylogenetics. Oxford University Press
- Nilsson K, Lindquist O, Pålsson C (1999) Association of *Rickettsia helvetica* with chronic perimyocarditis in sudden cardiac death. *Lancet* 354:1169–1173. [https://doi.org/10.1016/S0140-6736\(99\)04093-3](https://doi.org/10.1016/S0140-6736(99)04093-3)
- Nilsson K, Elfving K, Pålsson C (2010) *Rickettsia helvetica* in patient with meningitis, Sweden, 2006. *Emerg Infect Dis* 16:490–492. <https://doi.org/10.3201/eid1603.090184>
- Ogden NH, Lindsay LR (2016) Effects of climate and climate change on vectors and vector-borne diseases: ticks are different. *Trends Parasitol* 32:646–656. <https://doi.org/10.1016/j.pt.2016.04.015>
- Orkun Ö, Karaer Z, Çakmak A, Nalbantoğlu S (2014) Spotted fever group rickettsiae in ticks in Turkey. *Ticks Tick Borne Dis* 5:213–218. <https://doi.org/10.1016/j.ttbdis.2012.11.018>
- Otranto D (2018) Arthropod-borne pathogens of dogs and cats: from pathways and times of transmission to disease control. *Vet Parasitol* 251:68–77. <https://doi.org/10.1016/j.vetpar.2017.12.021>
- Otranto D, Dantas-Torres F, Breitschwerdt EB (2009a) Managing canine vector-borne diseases of zoonotic concern: part one. *Trends Parasitol* 25:157–163. <https://doi.org/10.1016/j.pt.2009.01.003>
- Otranto D, Dantas-Torres F, Breitschwerdt EB (2009b) Managing canine vector-borne diseases of zoonotic concern: part two. *Trends Parasitol* 25:228–235. <https://doi.org/10.1016/j.pt.2009.02.005>
- Otranto D, Paradies P, De Caprariis D et al (2009c) Toward diagnosing *Leishmania infantum* infection in asymptomatic dogs in an area where leishmaniasis is endemic. *Clin Vaccine Immunol* 16:337–343. <https://doi.org/10.1128/CVI.00268-08>
- Otranto D, Dantas-Torres F, Mihalca AD et al (2017) Zoonotic parasites of sheltered and stray dogs in the era of the global economic and political crisis. *Trends Parasitol* 33:813–825. <https://doi.org/10.1016/j.pt.2017.05.013>
- Otranto D, Iatta R, Baneth G et al (2019) High prevalence of vector-borne pathogens in domestic and wild carnivores in Iraq. *Acta Trop* 197: 105058. <https://doi.org/10.1016/j.actatropica.2019.105058>
- Panarese R, Iatta R, Latrofa MS et al (2020) Hyperendemic *Dirofilaria immitis* infection in a sheltered dog population: an expanding threat in the Mediterranean region. *Int J Parasitol* 50:555–559. <https://doi.org/10.1016/j.ijpara.2020.04.002>
- Parola P, Paddock CD, Socolovschi C et al (2013) Update on tick-borne rickettsioses around the world: a geographic approach. *Clin Microbiol Rev* 26:657–702. <https://doi.org/10.1128/CMR.00032-13>
- Regnery RL, Spruill CL, Plikaytis BD (1991) Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J Bacteriol* 173:1576–1589. <https://doi.org/10.1128/jb.173.5.1576-1589.1991>
- Rezaei A, Gharibi D, Pourmahdi Borujeni M, Mosallanejad B (2016) Seroprevalence of Lyme disease and Q fever in referred dogs to veterinary hospital of Ahvaz. *Iran Vet J* 11:34–41. <https://doi.org/10.22055/IVJ.2016.13016>
- Rubini AS, Paduan KS, Martins TF et al (2009) Acquisition and transmission of *Hepatozoon canis* (Apicomplexa: Hepatozoidae) by the tick *Amblyomma ovale* (Acari: Ixodidae). *Vet Parasitol* 164:324–327. <https://doi.org/10.1016/j.vetpar.2009.05.009>
- Shokri A, Fakhar M, Teshnizi SH (2017) Canine visceral leishmaniasis in Iran: a systematic review and meta-analysis. *Acta Trop* 165:76–89. <https://doi.org/10.1016/j.actatropica.2016.08.020>
- Soltani R, Dalimi A (2018) A molecular study on *Hepatozoon canis* infection in dogs in Tehran (Iran). *Arch Razi Inst* 73:257–263. <https://doi.org/10.22092/ARI.2017.110293.1125>
- Tabar M-D, Altet L, Francino O et al (2008) Vector-borne infections in cats: molecular study in Barcelona area (Spain). *Vet Parasitol* 151: 332–336. <https://doi.org/10.1016/j.vetpar.2007.10.019>
- Tamura K (1992) Estimation of the number of nucleotide substitutions when there are strong transition-transversion and G+C-content biases. *Mol Biol Evol* 9:678–687. <https://doi.org/10.1093/oxfordjournals.molbev.a040752>
- Travi BL, Cordeiro-da-Silva A, Dantas-Torres F, Miró G (2018) Canine visceral leishmaniasis: diagnosis and management of the reservoir living among us. *PLoS Negl Trop Dis* 12:1–13. <https://doi.org/10.1371/journal.pntd.0006082>
- Ullah Q, El-Adawy H, Jamil T et al (2019) Serological and molecular investigation of *Coxiella burnetii* in small ruminants and ticks in Punjab, Pakistan. *Int J Environ Res Public Health* 16:4271. <https://doi.org/10.3390/ijerph16214271>
- Yaghoobi-Ershadi MR (2016) Control of phlebotomine sand flies in Iran: a review article. *J Arthropod Borne Dis* 10:429–444

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.