

The effect of Diode Laser on the Candida Albicans Colonization in complete denture wearers

FATEMEH OWLIA¹, MOHAMMAD HASAN AKHAVAN KARBASSI², MOHAMMAD NIMA MOTALLAEI^{3*}

¹Department of oral and maxillofacial medicine, School of dentistry, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

²Department of oral and maxillofacial medicine, School of dentistry, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

³Department of oral and maxillofacial surgery, School of dentistry, Tehran University of Medical Sciences, Tehran, Iran.

Correspondence to Mohammad Nima Motallaei. Email: nimamotallaei@gmail.com, Tel: +98 9120620779,

ABSTRACT

Aim: To investigate the effect of diode laser on the Candida albicans colonization in complete denture patients.

Methods: A total of 40 complete maxillary dentures wearers studied in this cross-sectional study. Dentures were then randomly divided into two groups. In first group as the control group no intervention was made. Dentures of the second group merely exposed to laser irradiation by the diode laser with wavelength of 940 nm with 100mW for 30 seconds on mucosal surface of dentures before delivery. Samples were collected from the mucosal surface of upper dentures on the determined days. All collected colonies were counted. Clinical outcome was evaluated regarding the colonization of Candida albicans reported by CFU. Difference of colony counts results between two groups was evaluated by Mann-Whitney test. Statistical significance was assumed if $p < 0.05$.

Results: The results showed laser irradiation was significantly effective in reducing candida colonization. The lower density of the colonies was seen in laser group in compare with control group on 15th and 60th days after denture delivery ($P=0.002$) & ($P=0.003$), respectively.

Conclusions: generalization our data into the clinical setting, it can be proposed that a direct laser-based approach can significantly reduce the colonization of Candida albicans. It can also be concluded that laser light at specific wavelengths could be a possible promising novel strategy for prevention of denture stomatitis.

Keywords: Candida albicans; Stomatitis; Denture, complete; laser, Diode

INTRODUCTION

Increase in aging population is related to longer life expectancy^{1, 2}. The update rates of edentulism have been estimated up to 69 percent of the adult population in the world³. One of the main demand of the old is removable denture¹. although new methods for tooth replacement were developed, complete denture wearers are a large group of geriatrics, yet⁴⁻⁷. Regarding to rise of immunocompromised patients, such as those receiving chemotherapy or cytotoxic drugs, fungal infections have increased in recent years⁸⁻¹². One of the most common oral mucosal discomforts in denture wearers is denture stomatitis¹. Candida albicans has the most important role in the development of denture stomatitis. For sure the presence of this fungus in the mouth is regarded as commensalism, the transition from normal flora to pathogen organisms is related to an imbalance that occurs between the host and the fungus. This percent will be increased to 60-100% in denture wearers¹³⁻¹⁷. 24 hours use of prostheses, trauma to the tissues due to inadequate occlusion, poor oral hygiene, endocrine and immunological disorders, xerostomia, different type of immunodeficiency, allergic reaction to dental materials, trauma and antibiotic drugs and malignant diseases are some of the predisposing factors. The main underlying factor for denture stomatitis is denture itself¹⁸.

Candida albicans could stick to both mucosal and denture surface as an effective step to occurrence of denture stomatitis. Treatment of denture stomatitis can be a challenge due to its complicated etiology. Many efforts have been made to use different methods to manage this lesion from oral hygiene promotion, topical and systemic antifungal agents to denture replacement^{18, 19, 20, 21}. Many efforts have been made to use different methods to prevent

or treat this lesion. The use of Denture liners containing antifungal agents is a helpful method for the treatment of denture stomatitis in the short time. Use of optical sensitizers in combination with photodynamic therapy in some patients was effective^{22, 23}. Also, the use of microwaves has been effective for its management¹⁸. Diode laser whose effects are based on the decreased prosity of tissue surface of denture have recently been considered for the prevention of candida colonization^{18, 24, 25}. Other studies paid more attention to treatment of denture stomatitis than prevention^{18, 19, 22, 24, 26}. Basso FG et al claimed that LLLT had an inhibitory effect on the microorganisms, and this capacity can be altered according to the interactions between different microbial species²⁷. The role of diode laser on treatment of candidiasis has recently been taken into account. This study aimed to investigate the effect of diode laser on the Candida albicans colonization in complete denture wearers.

METHODS

Study design: This study was performed under the precepts of the World Medical Association's Declaration of Helsinki, as adopted in 2013 in Brazil. Total of 40 complete maxillary dentures from patients who were referred the Prosthetic Department of Yazd dental faculty, were consecutively examined and whom met inclusion criteria were selected. They randomly divided to two groups by random number table. A complete medical and dental history was taken from all of the participants.

Inclusion and exclusion criteria: Medical and dental history of all participants were obtained. No participant in either group had received antibiotics, steroids or immune therapy or used any antiseptic mouthwash during the study. Subjects, who reported history of any oral and

systemic diseases or taking any drugs or having previous denture, were excluded from this study.

Sampling: In laser group, tissue surface of dentures was irradiated by the diode laser (A.R.C GmbH Germany) with a wavelength of 940 nm. The laser was operated at a continuous mode at a distance of 5–10 mm perpendicular to the surface. The laser was used for 30 seconds with air- and water-cooling spray, and the average output power was 0.1 W. In control group no intervention was made. Samples were obtained by swabbing from the palatal impression surface of upper dentures before delivery & on the 15 & 60 days after repeating irradiation with the same process, and cultured on Sabouraud glucose agar plates (Figure 1). All collected yeasts were first counted and then identified by sub-culturing on CHROM agar Candida (CHRO-Magar, France)²⁸. Meanwhile, both groups received the same training for denture health. The microscopic morphology was investigated by counting the number of colony counts after being in the incubator for 48 hours.

Data analysis: Data were fed into SPSS-17 software. To analyze the normality of data, Kolmogorov-Smirnov and Shapiro-Wilk tests were used. If data were not normally distributed, the Kruskal-Wallis test was run to analyze the mean differences. Then, the Mann-Whitney test was used to analyze the differences between the means of collected colonies. The level of significance for pair comparisons was considered as $P < 0.05$.

Ethical consideration: Ethical approval was obtained from the research ethics committee of Shahid Sadoughi University of medical sciences. All the patients signed an informed consent form before the initiation of research.

RESULTS

Thirty three out of the 40 complete denture patients continued the study, 20 patients in the control group and 13 patients in the laser group. The means and standard deviation of two study groups are presented in Table 1. The laser group composed of 9 men and 4 women and with mean and standard deviation of age was 55.84 ± 11.52 years. The age range was between 41 to 78 years old. The control group included 20 persons (12 men and 8 women). The age range of patients was 32 to 78 and mean age of 56.35 ± 13.22 years. There was not any significant age difference between two groups.

Positive culture of Candida enrolled on the denture of all subjects on the 15 & 60 days after denture delivery meanwhile the average of colony count in both groups was near to zero on the day of delivery. The samples were reported above 1000, converted to 1000. (Table 1)

The colony count difference between two groups was significant (p -value = 0.003) on 15th. On the 60th day after delivery one patient did not come back for the culture therefore he was omitted. The difference in both groups was significant on 60th days after delivery. (P -value = 0.002)

After primary analysis, the data were assumed as quantitative by chi square test and cut of point of 500 (Table 2). As it is shown in the table, there was no significant difference between two groups on 15th day. (P value = 0.438)

There was a significant difference between the groups on the sixty days after irradiation. (P -value=0.007)(Table 3).

Table 1: Comparison of the average number of colony counts in two groups on the 15th & 60th.

Groups at different times	n	Mean and standard deviation	P value
Laser 15th	13	370/88+168/08	0/003
Control 15th	20	435/58+357/05	
Laser 60th	13	370/3+61/174	0/002
Control 60th	19	470/12+575/45	

Table 2: Frequency distribution of Candida colony count on the 15th day after laser irradiation

Colony count	Laser		Control	
	n	n	n	%age
<500	11	84/6	14	70
≤500	2	15/4	6	30
Total	13	100	20	100


P value = 0/438

Table 3: Frequency distribution of candidiasis colony count on the 60 days after laser irradiation

Colony count	Laser		Control	
	n	n	n	%age
<500	11	84/6	7	36/84
≤500	2	15/4	12	63/15
Total	13	100	19	100

P -value=0.007

Figure-1: Candida cultured on Sabouraud glucose agar plates



DISCUSSION

One of the most important concerns of dentists is the incidence and development of denture stomatitis. Numerous methods have been proposed for the prevention and treatment of this lesion. The use of laser irradiation, cryosurgery, electro surgery, and blade surgery have been successful results^{9, 22, 25, 29-32}. In some cases, in addition to improve oral health old denture should be replaced. Palatal surface of denture could act as a shelter for candida albicans than other surfaces. Use of disinfectant on denture surface is recommended in treatment of denture stomatitis. Microwave radiation has been shown to be effective disinfectant but hardness of the acrylic surfaces maybe impacted. Not only could laser irradiation be used for treatment of candidiasis, but also as a preventive method^{24, 25, 33-36}.

In this study, the mucosal surfaces of dentures were irradiated with a wavelength of 940 nm. It was operated at a continuous mode at a distance of 5–10 mm perpendicular to the surface before delivery. In some literatures both

tissue surface of the denture and palate were irradiated^{24, 34, 35}. In other studies only the palatal surfaces were affected, while in the present study, only palatal surface of dentures were irradiated^{33, 36, 37}. It is worthy to be mentioned, No exposure to participants, doing all steps of study in laboratory & prevention of multiple visits were some advantages of this method rather than others^{24, 34, 35}. In this study, a diode laser with wavelength of 940 nm with a power of 0.1W for 30 seconds was used but in Sivakumar study, the diode laser (wavelength of 980 nm & power of 2 W) irradiated two times for 15 seconds per visit & treatment procedure was completed in two visits. It was used two times in each sessions³⁶. In Scwinger study wavelength of 660nm of diode laser with a power of 30mw and the intensity of 7.5 J / cm² was used in mucosal contact for 10 seconds³⁷. In the Biscanin study, two diode lasers with a wavelength of 830 nm with intensity of 3 J / cm² for 60 minutes and another 685 nm with a power of 30 MW and 3 J / cm² for 10 minutes were used²⁵. Alves, a 660 nm diode laser with an intensity of 50 J / cm² for 20 minutes used in conjunction with a Photodithazine optical sensitizer³⁴. In Fontes study, a laser with a wavelength of 660 nm 100 MW and energy 4 Jules per cm²³⁵. In the Abduljabbar study, a LED with a wavelength of 440 to 460 nm with a power of 24 MW / cm² for a denture and a 260 MW LED and an intensity of 102 MW / cm² was applied on tissue surface³³. Almost the wavelength & duration of laser have been used in the present study were similar to Sivakumar³⁶. It led to minimum damage to denture in removing porosity.

The samples were taken with sterilized swabs and immediately cultured in a dextrose agar. This work has been done before denture delivery and was repeated on 15th and 60th days later. The colony count was calculated by the relevant expert after 48 hours remaining in the incubator. Some studies used the diode laser for treatment of candidiasis. Quite the opposite, there is no report about the use of diode laser for the prevention of candida colonization. Sivakumar used exfoliative cytology and PAS dyeing only before denture stomatitis³⁶. Scwinger sampled before delivery to the patients, immediately after delivery and on 7th, 15th, and 30 days later³⁷. Biscanin sampled from denture and palate before and after laser irradiation but Alves study on 15th, 30, and 45 days after that.²⁵,³⁴ Abduljabbar carried out exfoliative cytology, PAS staining, sampling and culture before delivery and 3 months later³³. In order to ensure no contamination of the denture, culture before delivery was mandatory. The colony count before delivery was not zero in two cases, which was related to contamination, probably in the laboratory, but the statistical difference was not significant. Of course, the risk of candidiasis after delivery increased over time due to mucosal contact. One of the person in control group was missed, because she did not come back for third step of culture. The least time for primary and secondary candida colonization on palatal surface of denture is 15th & 60th days with p-value=0.003 & p-value=0.002 respectively, that were selected in the study^{34, 37}.

In statistical analysis at zero time, there was no significant difference between groups, in other words they were similar (pvalue>0.05). Assuming the quantitative value of the data, the difference between the two groups was

significant at 15th days after delivery (p = 0.003). Considering data qualification. Analyzing with chi-square & non parametric test, It was concluded that difference was statistically significant only on the 60th day (p-value=0.007). Little sample size was the main limitation of this study that should be considered in future studies.

CONCLUSION

Mycological findings in the present study revealed that application of Diode laser with this process could prevent candida colonization in long term (60 days) more than short term (15 days). By elimination of micro porosity of denture and improving their quality, Candida colonization and probability of denture stomatitis could be decreased.

Financial disclosure: This study has been supported financially by the Vice Chancellor for Research and Technology of Yazd Shahid Sadoughi University of Medical Sciences.

Conflict of interest: There are no conflicts of interest.

REFERENCES

- Gleiznys A, Zdanaviciene E, Zilinskas J. Candida albicans importance to denture wearers. A literature review. *Stomatologija*. 2015;17(2):54-66.
- Sanderson WC, Scherbov S. Faster increases in human life expectancy could lead to slower population aging. *PLoS one*. 2015;10(4):e0121922.
- Wu B, Liang J, Landerman L, Plassman B. Trends of edentulism among middle-aged and older Asian Americans. *American journal of public health*. 2013 Sep;103(9):e76-82.
- Nitecka-Buchta A, Proba T, Proba P, Stefanski K, Baron S. Functional Assessment of the Stomatognathic System, after the Treatment of Edentulous Patients, with Different Methods of Establishing the Centric Relation. *Pain research & management*. 2018;2018:1572037.
- Mosaddad SA, Beigi K, Doroodizadeh T, Haghnegahdar M, Golfeshan F, Ranjbar R, et al. Therapeutic applications of herbal/synthetic/bio-drug in oral cancer: An update. *Eur J Pharmacol*. 2021 2021/01/05;890:173657.
- Yazdani M, Tabesh H, Houshmand B, Tebyanian H, Soufdoost RS, Tahmasebi E, et al. Fabrication and properties of β TCP/Zeolite/Gelatin scaffold as developed scaffold in bone regeneration: in vitro and in vivo studies. *Biocyberrn Biomed Eng*. 2020 2020/10/01;40(4):1626-37.
- Yazdani M, Rahmani A, Tahmasebi E, Tebyanian H, Yazdani A, Mosaddad SA. Current and advanced nanomaterials in dentistry as regeneration agents: an update. *Mini-Rev Med Chem*. 2020;20:1-21.
- Garbee DD, Pierce SS, Manning J. Opportunistic Fungal Infections in Critical Care Units. *Critical care nursing clinics of North America*. 2017 Mar;29(1):67-79.
- Moghadam ET, Yazdani M, Tahmasebi E, Tebyanian H, Ranjbar R, Yazdani A, et al. Current herbal medicine as an alternative treatment in dentistry: In vitro, in vivo and clinical studies. *Eur J Pharmacol*. 2020 2020/12/15;889:173665.
- Zolfaghar M, Amoozegar MA, Khajeh K, Babavalian H, Tebyanian H. Isolation and screening of extracellular anticancer enzymes from halophilic and halotolerant bacteria from different saline environments in Iran. *Mol Biol Rep*. 2019;46(3):3275-86.
- Mosaddad SA, Tahmasebi E, Yazdani A, Rezvani MB, Seifalian A, Yazdani M, et al. Oral microbial biofilms: an update. *Eur J Clin Microbiol*. 2019 2019/08/01;38:2005-19.
- Rezaeeyan Z, Safarpour A, Amoozegar MA, Babavalian H, Tebyanian H, Shakeri F. High carotenoid production by a halotolerant bacterium, *Kocuria* sp. strain QWT-12 and anticancer activity of its carotenoid. *EXCLI J*. 2017;16:840-51.

13. Chopde N, Jawale B, Pharande A, Chaudhari L, Hiremath V, Redasani R. Microbial colonization and their relation with potential cofactors in patients with denture stomatitis. *The journal of contemporary dental practice.* 2012 Jul 1;13(4):456-9.
14. Loster BW, Loster J, Wieczorek A, Ryniewicz W. Mycological analysis of the oral cavity of patients using acrylic removable dentures. *Gastroenterology research and practice.* 2012;2012:951572.
15. Sampaio-Maia B, Figueiral MH, Sousa-Rodrigues P, Fernandes MH, Scully C. The effect of denture adhesives on *Candida albicans* growth in vitro. *Gerodontology.* 2012 Jun;29(2):e348-56.
16. Seifi Kafshgari H, Yazdani M, Ranjbar R, Tahmasebi E, Mirsaeed S, Tebyanian H, et al. The effect of *Citrullus colocynthis* extracts on *Streptococcus mutans*, *Candida albicans*, normal gingival fibroblast and breast cancer cells. *J Biol Res.* 2019;92(1):8201.
17. Heidari MF, Arab SS, Noroozi-Aghideh A, Tebyanian H, Latifi AM. Evaluation of the substitutions in 212, 342 and 215 amino acid positions in binding site of organophosphorus acid anhydrolase using the molecular docking and laboratory analysis. *Bratisl Lek Listy.* 2019;120(2):139-43.
18. Neppelenbroek KH, Pavarina AC, Palomari Spolidorio DM, Sgavioli Massucato EM, Spolidorio LC, Vergani CE. Effectiveness of microwave disinfection of complete dentures on the treatment of *Candida*-related denture stomatitis. *Journal of oral rehabilitation.* 2008 Nov;35(11):836-46.
19. Souza RC, Junqueira JC, Rossoni RD, Pereira CA, Munin E, Jorge AO. Comparison of the photodynamic fungicidal efficacy of methylene blue, toluidine blue, malachite green and low-power laser irradiation alone against *Candida albicans*. *Lasers in medical science.* 2010 May;25(3):385-9.
20. Ribeiro Rocha GDS, Neves Duarte T, de Oliveira Corrêa G, Nampo FK, de Paula Ramos S. Chemical cleaning methods for prostheses colonized by *Candida* spp.: A systematic review. *The Journal of prosthetic dentistry.* 2020 Dec;124(6):653-8.
21. Sugio CYC, Garcia A, Albach T, Moraes GS, Bonfante EA, Urban VM, et al. *Candida*-Associated Denture Stomatitis and Murine Models: What Is the Importance and Scientific Evidence? *Journal of fungi (Basel, Switzerland).* 2020 May 23;6(2).
22. Marei MK, Abdel-Meguid SH, Mokhtar SA, Rizk SA. Effect of low-energy laser application in the treatment of denture-induced mucosal lesions. *The Journal of prosthetic dentistry.* 1997 Mar;77(3):256-64.
23. Wilson M, Mia N. Sensitisation of *Candida albicans* to killing by low-power laser light. *Journal of oral pathology & medicine : official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology.* 1993 Sep;22(8):354-7.
24. Maver-Biscanin M, Mravak-Stipetic M, Jerolimov V. Effect of low-level laser therapy on *Candida albicans* growth in patients with denture stomatitis. *Photomedicine and laser surgery.* 2005 Jun;23(3):328-32.
25. Maver-Biscanin M, Mravak-Stipetic M, Jerolimov V, Biscanin A. Fungicidal effect of diode laser irradiation in patients with denture stomatitis. *Lasers in surgery and medicine.* 2004;35(4):259-62.
26. Martins KV, de Lacerda Gontijo SM. Treatment of denture stomatitis: literature review. *Revista Brasileira de Odontologia.* 2017;74(3):215-20.
27. Basso FG, Oliveira CF, Fontana A, Kurachi C, Bagnato VS, Spolidorio DM, et al. In Vitro effect of low-level laser therapy on typical oral microbial biofilms. *Brazilian dental journal.* 2011;22(6):502-10.
28. Lotfi-Kamran MH, Jafari AA, Falah-Tafti A, Tavakoli E, Falahzadeh MH. *Candida* Colonization on the Denture of Diabetic and Non-diabetic Patients. *Dental research journal.* 2009 Spring;6(1):23-7.
29. Tahmasebi E, Alikhani M, Yazdani A, Yazdani M, Tebyanian H, Seifalian A. The current markers of cancer stem cell in oral cancers. *Life Sci.* 2020 2020/05/15;249:117483.
30. Tahmasebi E, Alam M, Yazdani M, Tebyanian H, Yazdani A, Seifalian A, et al. Current biocompatible materials in oral regeneration: a comprehensive overview of composite materials. *J Mater Res Technol.* 2020 2020/09/01;9(5):11731-55.
31. Mosaddad SA, Yazdani M, Tebyanian H, Tahmasebi E, Yazdani A, Seifalian A, et al. Fabrication and properties of developed collagen/strontium-doped Bioglass scaffolds for bone tissue engineering. *J Mater Res Technol.* 2020 2020/11/01;9(6):14799-817.
32. Soufdoost RS, Yazdani M, Tahmasebi E, Yazdani A, Tebyanian H, Karami A, et al. In vitro and in vivo evaluation of novel Tadalafil/ β -TCP/Collagen scaffold for bone regeneration: A rabbit critical-size calvarial defect study. *Biocybern Biomed Eng.* 2019 2019/07/20;39(3):789-96.
33. Abduljabbar T, Al-Askar M, Baig MK, AlSowaygh ZH, Kellesarian SV, Vohra F. Efficacy of photodynamic therapy in the inactivation of oral fungal colonization among cigarette smokers and non-smokers with denture stomatitis. *Photodiagnosis and photodynamic therapy.* 2017 Jun;18:50-3.
34. Alves F, Alonso GC, Carmello JC, Mima EGO, Bagnato VS, Pavarina AC. Antimicrobial Photodynamic Therapy mediated by Photodithazine((R)) in the treatment of denture stomatitis: A case report. *Photodiagnosis and photodynamic therapy.* 2018 Mar;21:168-71.
35. Da Costa Fontes KBF, Dos Santos Farias IB, Cappato LP, Da Silva BA, De Souza Azevedo R, Tucci R, et al. Op - *Candida*-Associated Denture Stomatitis Treated with Antimicrobial Photodynamic Therapy: Four Cases. *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology.* 2017;123(2):e34.
36. Sivakumar K, Palanivelu S, Ramalingam S, Seshadri S. Denture Stomatitis: Treatment with Diode Laser. *International Journal of Prosthodontics and Restorative Dentistry.* 2011;1(1):55-7.
37. Scwingel AR, Barcessat AR, Nunez SC, Ribeiro MS. Antimicrobial photodynamic therapy in the treatment of oral candidiasis in HIV-infected patients. *Photomedicine and laser surgery.* 2012 Aug;30(8):429-32