

## Evaluation of In Vitro Effectiveness of Seven Disinfectants over Controlling *Candida* on Complete Dentures

### Dear Editor,

Despite major advances in dentistry and dental prostheses, complete removable dentures are still widely used for oral rehabilitation of edentulous people. Denture surfaces prepare a favorable environment for adherence and proliferation of oral endogenous microorganisms, which is a crucial first step in the initiation of biofilm formation, caused denture stomatitis.<sup>1</sup> *Candida* species are known as the predominant oral etiologic agent of denture stomatitis, a common mucosal inflammation usually seen on the denture-bearing mucosa of denture wearers.<sup>2,3</sup> There are several physical and chemical methods for cleaning of denture plaques.<sup>4</sup> Overnight soaking of denture in chemical solutions' including denture cleansers and detergents is an applicable and inexpensive method for improving of denture hygiene.<sup>5</sup> Ideal denture cleansers and disinfectants should be bactericidal, fungicidal, nontoxic and harmless to denture's structure, effective for removing of organic and inorganic deposits on the denture.<sup>4</sup>

To compare the effectiveness of seven commercially available disinfectants for cleaning of complete denture, we performed an experimental study in Yazd Dentistry School. Sixty three complete maxillary dentures, which were used more than one year, were randomly divided to nine groups. Cultures from their internal surface were obtained by vigorous rubbing with a sterile cotton swab onto Sabouraud dextrose agar plates, incubated 48 hours at 30°C for quantification of isolated *Candida* colonies. Isolated *Candida* colonies were also identified using germ tube test. Denture were then immersed for 1 hour in sterile 'zip-lock' bags filled with 200 ml of sodium hypochlorite (2%, 1% and 0.5%), 0.2% chlorhexidine, 4% benzalconium chloride, 1% deconex and normal saline. Sterile distilled water and 100,000 IU solution of nystatin were also used as negative and positive controls in the current study. After the disinfectant protocol, the dentures were washed with sterile distilled water, re-swabbed and cultured as previously described before the disinfection. All cultures were performed by a single operator using sterile gloves, which were dis-

posed and replaced for each denture. Density of isolated *Candida* colonies before and after disinfection was compared with Wilcoxon statistical (<0.05) test using SPSS software.

All dentures were positive for *Candida* species in the initial culture before disinfection protocol (Table 1). *Candida albicans* was the most common isolated *Candida* species isolated from 71% of dentures, followed by non-*albicans Candida* species, isolated from 23% of dentures and 6% had mix species. All immersion solutions were found to reduce the growth of *Candida* in comparing with the initial culture. Two percent sodium hypochlorite ( $p=0.008$ ), 1% sodium hypochlorite ( $p=0.016$ ), 0.2% chlorhexidine ( $p=0.025$ ) and 1% deconex ( $p=0.046$ ) were known as the most effective disinfectants shown the maximum reduction in the load of isolated *Candida* species in current study (Table 2).

**Table 1:** Positive cultures of *Candida* on dentures before and after protocol, the numbers indicate the positive growth from total of seven dentures

Disinfectant solutions	Before	After
2% Sodium hypochlorite	7	0
1% Sodium hypochlorite	7	1
0.5% Sodium hypochlorite	7	1
0.2% Chlorhexidine	7	2
1% Deconex	7	2
4% Benzalconium chloride	7	3
Normal saline	7	4
Nystatine	7	0
Distilled water	7	7

The results of the current study showed that all the initial cultures obtained from dentures, which taken directly from the patients' mouth were positive for *Candida* species as expected. The objective of denture disinfection by immersing in chemicals is to obtain a clean, decontaminated prosthesis by removing the oral microbial contamination.<sup>5</sup> There are many studies, which showed more efficacy of chemical than physical methods.<sup>6</sup> The 2% and 1% sodium hypochlorite showed the highest cleaning effect followed by 0.2% chlorhexidine in the current study as also reported by Montagner *et al.*,<sup>7</sup> Furthermore 0.5%

**Table 2:** Load of *Candida* colonies (CFU\*) on the dentures

Disinfectant solutions	Initial culture Mean (SD)	After the protocol Mean (SD)	P value
2% Sodium hypochlorite	202 (45.2)	3 (1)	0.008
1% Sodium hypochlorite	75.3 (34.8)	4.1 (2)	0.016
0.5% Sodium hypochlorite	119 (39.5)	6.2 (2.3)	0.036
0.2% Chlorhexidine	152 (48.3)	5.5 (2)	0.025
1% Deconex	89.2 (27)	11 (3)	0.046
4% Benzalconium chloride	109.3 (51.2)	15 (3.2)	0.066
Normal saline	121.7 (32.9)	29 (5.7)	0.083
Nystatine	155 (51.2)	0	0.0001
Distilled water	68.5 (8)	43.2 (18)	0.45

\*Colony forming unit in each culture

sodium hypochlorite, deconex (1%) and 4% benzalconium chloride showed a lower effect in disinfection of denture in the current study. Yilmaz *et al.* used 5.25% and 2% sodium hypochlorite as well as deconex and salvex and showed the same results;<sup>8</sup> however 5% sodium hypochlorite can damage dentures<sup>9</sup> and was not used in the present study.

**Keywords:** In vitro; *Candida*; Disinfectant; Denture

**Conflict of interest:** None declared.

**AA Jafari<sup>1</sup>, A Falah-Tafti<sup>2\*</sup>, MH Lotfi-**

**Kamran<sup>2</sup>, H Fallahzadeh<sup>3</sup>, F Akaberi<sup>4</sup>**

<sup>1</sup>Department of Parasitology and Mycology, Medical School, <sup>2</sup>Department of Prosthodontics, Dentistry School, <sup>3</sup>Department of Biostatistics and Epidemiology, School of Public Health, <sup>4</sup>Dentist, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

\*Correspondence: Abbas Falah Tafti, PhD, Department of Prosthodontics, Dentistry School, Daha Fajr Blvd, Shahid Sadoughi University of Medical Sciences, Yazd, Iran. Tel: +98-913-3519212, e-mail: jaabno@gmail.com  
Received: August 20, 2011 Accepted: November 22, 2011

## References

- Webb BC, Thomas CJ, Willcox MD, Harty DW, Knox KW. Candida-associated denture stomatitis. Aetiology and management: a review. Part 1. Factors influencing distribution of *Candida* species in the oral cavity. *Aust Dent J* 1998;**43**:45-50. [9583226] [http://dx.doi.org/10.1111/j.1834-7819.1998.tb00152.x]
- Barbeau J, Séguin J, Goulet JP, de Koninck L, Avon SL, Lalonde B, Rompré P, Deslauriers N. Reassessing the presence of *Candida albicans* in denture-related stomatitis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2003;**95**:51-9. [1253902] [http://dx.doi.org/10.1067/moe.2003.44]
- Gendreau L, Loewy ZG. Epidemiology and Etiology of Denture Stomatitis. *J Prosthodont* 2011;**20**: 251-60. [21463383] [http://dx.doi.org/10.1111/j.1532-849X.2011.00698.x]
- Abelson DC. Denture plaque and denture cleansers: review of the literature. *Gerodontology* 1985;**1**:202-6. [3005102]
- Pavarina ac, Pizzolitto AC, Machado AI, Vergani CE, Giampaolo ET. An infection control protocol: effectiveness of immersion solutions to reduce the microbial growth on dental prostheses. *J Oral Rehabil* 2003;**30**: 532-6. [12752936] [http://dx.doi.org/10.1046/j.1365-2842.2003.01093.x]
- Buergers R, Rosentritt M, Schneider-Brachert W, Behr M, Handel G, Hahnel S. Efficacy of denture disinfection methods in controlling *Candida albicans* colonization in vitro. *Acta Odontol Scand* 2008;**66**:174-80. [18568477] [http://dx.doi.org/10.1080/00016350802165614]
- Montagner H, Montagner F, Braun KO, Peres PE, Gomes BP. In vitro antifungal action of different substances over microwaved-cured acrylic resins. *Appl Oral Sci* 2009;**17**:432-5. [19936521]
- Yilmaz H, Aydin C, Bal BT, Özçelik B. Effects of disinfectants on resilient denture-lining materials contaminated with *Staphylococcus aureus*, *Streptococcus sobrinus*, and *Candida albicans*. *Quintessence Int* 2005;**36**:373-81. [15892535]
- Falah-tafti A, Jafari AA, Lotfi-kamran MH. Comparison of the Effectiveness of Sodium Hypochlorite and Dentamize Tablet for Denture Disinfection. *World Journal of Medical Sciences* 2008;**3**:10-14.