



# Human sperm parameter improvement associated with *Ceratonia siliqua* extract as a cryopreservation supplement after vitrification

Tooba Farazmand<sup>1</sup>, Fatemeh Mansouri<sup>2</sup>, Yeganeh Koohestanidehaghi<sup>3</sup>, Erfan Shahandeh<sup>4</sup>

<sup>1</sup>Department of Gynecology, Faculty of Medicine, North Khorasan University of Medical Sciences, Bojnurd; <sup>2</sup>Department of Physiology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad; <sup>3</sup>Research and Clinical Center for Infertility, Yazd Reproductive Sciences Institute, Shahid Sadoughi University of Medical Sciences, Yazd; <sup>4</sup>Department of Biology, Islamic Azad University of Hamedan, Hamedan, Iran

**Objective:** Given the destructive effects of oxidative stress on sperm structure, this study was conducted to investigate the antioxidant effects of different concentrations of *Ceratonia siliqua* plant extract on human sperm parameters after the freezing-thawing process.

**Methods:** A total of 20 normozoospermic samples were frozen. Each sample was divided into two control groups (fresh and cryopreservation) and three cryopreservation experimental groups (containing *C. siliqua* extract at concentrations of 20, 30, and 40 µg/mL in the freezing extender). Motility, intracellular levels of reactive oxygen species (ROS), plasma membrane integrity (PMI), mitochondrial membrane potential (MMP), viability, and acrosome reaction parameters were evaluated.

**Results:** Statistical analysis showed that the highest motility, viability, and PMI were associated with the 20 µg/mL concentration of *C. siliqua* extract. At all concentrations, intracellular ROS levels were significantly lower and the levels of MMP and the acrosome reaction were significantly higher than in the cryopreservation control group ( $p \leq 0.05$ ).

**Conclusion:** *C. siliqua* extract supplements at concentrations of 20, 30, and 40 µg/mL improved sperm motility, viability, PMI, MMP, intracellular ROS, and the acrosome reaction.

**Keywords:** Antioxidants; Carob; Cryopreservation; Fabaceae; Sperm motility

## Introduction

Freezing human sperm is an effective and beneficial strategy in the field of male fertility [1]. This method can be used for men with cancer who are undergoing radiotherapy or chemotherapy treatment, in sperm donation programs, or in conjunction with surgical procedures that endanger male fertility [2-5]. The use of cryopreservation to create a sperm bank for healthy men who are exposed to ionizing radiation, biological contaminants, or toxins at work is an

other related goal [6]. Despite these benefits of sperm cryopreservation, we now understand that cryopreservation and thawing cause irrecoverable changes in sperm function and structure [7]. In the cryopreservation process, sperm are exposed to stressors such as osmotic pressure change, pH change, dehydration, the creation of ice crystals, and the generation of free radicals [8]. These factors can impair sperm motility, cell membrane and mitochondrial structure, chromatin structure, and sperm viability [9]. In recent years, extensive efforts have been made to improve cryopreservation and reduce the harmful effects of the freezing-thawing process on human sperm [10]. These include changes to cryopreservation methods, composition of the cryopreservation medium, cryopreservation and thawing times, and packaging of samples [11,12]. In general, freezing media include cryoprotectants, ionic or non-ionic materials to maintain the pH, and an energy substrate, as well as fatty acids, antibiotics, and antioxidants [11]. According to previous studies, the complementa-

Received: July 7, 2022 · Revised: December 20, 2022 · Accepted: January 14, 2023  
Corresponding author: Erfan Shahandeh

Department of biology, Islamic Azad University of Hamedan, Hamedan, Iran  
Tel: +98-09127276023 Fax: +98-09127276023 E-mail: esh957522@gmail.com

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

tion of freezing-thawing environments with various factors, including antioxidants, is an effective approach to improve the quality of frozen-thawed sperm [13]. Many studies have focused on plants with antioxidant capacities that can counteract the damaging effects of free radicals due to oxidative stress [14]. *Ceratonia siliqua* is an evergreen plant native to the Mediterranean region. *C. siliqua* contains compounds such as vitamins (B, C, D, and E), polyphenols, and minerals (iron, phosphorus, potassium, sodium, and calcium) [15,16]. The phenolic components of this plant improve oxidative stress conditions [17] and can act as a powerful source of antioxidants. As such, the use of phenolic antioxidants is recommended to improve oxidative damage, with minimal side effects and ease of use [12,18]. For the first time, in the current study, we simultaneously evaluated the effects of *C. siliqua* extract on human sperm parameters including motility, viability, cell membrane and mitochondrial potential, intracellular levels of reactive oxygen species (ROS), and the acrosome reaction using computer-assisted sperm analysis, fluorescence, hyposmotic swelling (HOS) testing, JC-1 staining, flow cytometry, and fluorescent thiocyanate, respectively.

## Methods

### 1. Extraction

#### 1) Preparation of *C. siliqua* hydroalcoholic extract

The *C. siliqua* plant was prepared by a medicinal plant expert from the local plant market and approved by a pharmacognosy expert. First, 100 g of fresh *C. siliqua* plant matter was dried at room temperature and then ground in a blender, yielding a homogeneous powder. Then, 10 g of *C. siliqua* powder was mixed with 1,000 mL of 96% ethanol (Merck). The resulting solution was added to distilled water in a 50:50 ratio and kept at room temperature for 72 hours. This extract was then filtered (filter paper no. 4; Whatman PLC). The filtered solution was concentrated at 50 °C using a rotary apparatus. Finally, a brown extract was obtained, dried in an oven at 40 °C, and kept at -20 °C until use [19].

### 2. Participation and semen collection

Verbal informed consent was obtained from each patient for the use of their semen samples (IR.ACECR.HAMEDAN.1400.140) were approved by the ethics committee of the Hamadan Branch of Azad University. A total of 20 normal sperm samples from men referred to the Aban Infertility Center between April 29, 2021 and August 17, 2021 were used. Samples were prepared during 3 to 5 days of abstinence. These normal samples had normal morphology (> 4%), normal motility (< 40%), and a concentration above 15 million/mL and were collected according to the 2010 World Health Organization guidelines.

### 3. Sperm processing

The swim-up method was used to process the semen samples [20]. In this method, the samples were first centrifuged for 5 minutes at 300 × g; then, the supernatant was removed, and the pellets were precipitated with preheated human tubal fluid (HTF) and 2.5% human serum albumin (HSA) (Vitrolife) [21]. These samples were poured into a Falcon tube and incubated at a 45° angle in an incubator (5% CO<sub>2</sub>, 37 °C) for 1 hour. The supernatant was poured into a microtube, analyzed, and then frozen by vitrification.

### 4. Cryopreservation and thawing

After the preparation of sperm, the samples were frozen via the microdroplet technique [22]. For the cryopreservation control group, the sperm solution and HTF were mixed with a solution containing 5% HSA and 0.5 mol/L sucrose (Vitrolife) at a 1:1 ratio. *C. siliqua* extract was prepared at several concentrations (20, 30, and 40 µg/mL). For the cryopreservation experimental groups, the sperm solution was mixed with HSA and sucrose solution along with 20, 30, or 40 µg/mL of *C. siliqua* extract.

Droplets of 30 µL of the micropipette-prepared suspensions were poured into a metal strainer and placed in liquid nitrogen, and the frozen samples were kept in nitrogen storage tanks for 1 week [23]. For the thawing process, we first incubated the HTF medium for 2 hours at 37 °C and then immersed the frozen samples in 5 mL of heated HTF supplemented with 1% HSA.

Next, the sperm suspensions were incubated at 37 °C and 5% CO<sub>2</sub> for 5 minutes. Finally, the samples were centrifuged for 5 minutes at 1,800 rpm, the pellets were resuspended in 50 µL HTF, and the sperm were evaluated [23].

### 5. Assessment of sperm parameters

#### 1) Motion characteristics

A computer-aided sperm analyzer system (Sperm Class Analyzer version 6; Microptic) was used to assess motility and the motility indices. In this study, 5-µL samples from each group were placed in a preheated Makler chamber (Proiser), and general motility, progressive movement and motor indices, average path velocity (µm/sec), curvilinear velocity (µm/sec), straight-line velocity (µm/sec), mean linearity (µm/sec), straightness (µm/sec), amplitude of lateral head displacement (µm), and beat cross frequency (Hz) were assessed for 500 sperm.

#### 2) Viability

To assess sperm viability, a LIVE/DEAD sperm kit was used according to the manufacturer's instructions (L-7011; Molecular Probes). To accomplish this, 0.1 µL of SYBR14 working solution (containing 150 mM sodium chloride, 10 mM HEPES, 10% bovine serum albumin,

and SYBR14) was added to 50  $\mu\text{L}$  of sperm suspension. The resulting solution was incubated at 37  $^{\circ}\text{C}$  for 10 minutes. Then, 1  $\mu\text{L}$  of propidium iodide solution was added to the previous suspension and incubated for another 5 minutes in an incubator at 37  $^{\circ}\text{C}$ , and 10  $\mu\text{L}$  of Hancock solution was added to the solution to immobilize the sperm. Finally, to assess viability, 6  $\mu\text{L}$  of this final suspension was placed on a glass slide, covered, and examined using a fluorescence microscope (CX21; Olympus; excitation at 450–490 nm, emission at 520 nm) with a magnification of  $\times 1,000$ . In this protocol, cells were stained with propidium iodide and SYBR14 on each slide. Dead and living spermatozoa were evaluated based on red and green color emissions, respectively. The LIVE/DEAD sperm ratio is depicted in Figures 1 and 2 [21].


### 3) Plasma membrane integrity

The HOS test was used to evaluate membrane integrity. We mixed 10  $\mu\text{L}$  of the sperm sample, 100  $\mu\text{L}$  of HOS solution containing 0.73 g of sodium citrate and 1.35 g of fructose (Merck), and 100 mL of distilled water and incubated the mixture (5%  $\text{CO}_2$ , 37  $^{\circ}\text{C}$ ). Then, 15  $\mu\text{L}$  of this solution was added to 5  $\mu\text{L}$  of eosin Y solution (2%), and a smear was prepared from the resulting mixture on a glass slide. Finally, 200 sperm were assessed using a light microscope (ECLIPSE


50i; Nikon) with a magnification of  $\times 1,000$ . Coiled-tail sperm were considered to have intact plasma membranes (Figure 3) [24].

### 4) Mitochondrial activity


JC-1 is a lipophilic cationic dye (T4069; Sigma-Aldrich) used to as-



**Figure 2.** The graph shows the ratio of live/death sperm in the control and experimental groups. The most positive effect on the ratio of live/death sperm is related to the group containing 20  $\mu\text{g/mL}$  cryopreservation supplement of siliqua extract. *C. siliqua*, *Ceratonia siliqua*.



**Figure 1.** This shows dead and live sperm based on the fluorescence emitted. Green sperm stained with SYBR14 are alive and orange sperm stained with propidium iodide are dead.



**Figure 3.** This figure shows the integrity of the plasma membrane based on hypoosmotic swelling tests. Sperms with straight tails are considered dead and sperms with coiled tails are considered alive.

sess sperm mitochondrial activity. For this assessment, sperm samples were centrifuged for 5 minutes at  $500 \times g$ , after which the pellet was diluted with phosphate-buffered saline. Then,  $500 \mu\text{L}$  of this solution was mixed with  $1 \mu\text{L}$  of JC-1 stock solution and incubated at  $37^\circ\text{C}$  for 40 minutes. Next,  $10 \mu\text{L}$  of Hancock solution was added to the previous solution to immobilize the sperm. Finally,  $2.5 \mu\text{L}$  of the prepared sample was placed on a glass slide, covered, and assessed under a BX51 fluorescence microscope (Olympus; excitation at  $450\text{--}490 \text{ nm}$ , emission at  $520 \text{ nm}$ ) at a magnification of  $\times 1,000$ , under which 200 sperm per slide were examined. Sperm with yellow/orange fluorescence at the midline were considered to display high mitochondrial activity, while sperm with green fluorescence were considered to exhibit low mitochondrial activity (Figure 4) [25].

##### 5) Acrosome integrity

To evaluate the integrity of the acrosome, fluorescein isothiocyanate-conjugated *Pisum sativum* agglutinin (FITC-PSA, L0770; Sigma-Aldrich) was used. Initially, a  $30\text{-}\mu\text{L}$  smear was prepared on a glass slide for each sample. After drying, the samples were fixed at room temperature with methanol for 30 minutes. Then,  $50 \mu\text{L}$  of


FITC-PSA solution was poured on each slide and incubated for 30 minutes. Finally, the stained slides were washed with distilled water and were evaluated using a BX51 fluorescence microscope (Olympus; excitation at  $450\text{--}490 \text{ nm}$ , emission at  $520 \text{ nm}$ ) with a magnification of  $\times 1,000$ . For each slide, 200 sperm were examined. Sperm with bright green fluorescence in the acrosome area were identified as having intact acrosomes, while sperm with no green or pale green fluorescence near the equator were identified as acrosome-reacted (Figure 5) [21].

##### 6) Intracellular ROS


To assess intracellular ROS, sperm samples were first suspended after melting with phosphate-buffered saline. Then,  $10 \mu\text{L}$  of dihydroethidium solution was added. The samples were incubated at room temperature for 30 minutes. For this measurement, flow cytometry was used. Red fluorescence with an FL2 detector ( $525\text{--}625 \text{ nm}$ ) was used to indicate intracellular ROS [26,27].

##### 7) Flowcytometric analysis

Flowcytometric analysis was conducted using Cyflogic version



**Figure 4.** This figure shows the activity of sperm mitochondria. Sperms with yellow fluorescence in the midpiece region show high activity of the mitochondrial membrane and sperms with green fluorescence show low activity of the mitochondrial membrane.



**Figure 5.** This figure shows the acrosome reaction based on the emission of different colors from the acrosome region using a fluorescence microscope. Sperms with clear green fluorescence as intact acrosomes and the sperms with pale green fluorescence were determined as acrosome reaction.

2.5.1 (CyFlo Ltd.). This analysis was performed using a FACSCalibur flow cytometer (BD Biosciences) and an argon laser with a wavelength of 488 nm [28].

## Results

As shown in Table 1, when either 30 or 40 µg/mL of *C. siliqua* extract was added to the sperm freezing extender, no significant differences in total or progressive motility were observed relative to the cryopreservation control group. In contrast, significantly greater motility was observed in the group treated with 20 µg/mL extract ( $p \leq 0.05$ ). No significant difference was observed in the other motility characteristics for any of the extract concentrations relative to the cryopreservation control group (Table 1).

As shown in Table 2, the intracellular ROS level was significantly reduced among sperm treated with any concentration of *C. siliqua* extract relative to the cryopreservation control group ( $p \leq 0.05$ ). Additionally, by increasing the concentration of *C. siliqua* extract from 20 µg/mL to 30 or 40 µg/mL, viability and plasma membrane integrity (PMI) were significantly improved in the cryopreservation experi-

mental groups relative to the cryopreservation control group. The greatest increase relative to the control was associated with the 20 µg/mL extract concentration, and no significant difference was observed between 30 and 40 µg/mL ( $p \geq 0.05$ ). Levels of both mitochondrial membrane potential (MMP) and the acrosome reaction showed a significant increase at all three concentrations compared to the cryopreservation control group. However, no significant difference was seen between the experimental groups (Table 2).

## Discussion

Sperm cryopreservation can involve the formation of ice crystals, osmotic imbalance, and oxidative stress, potentially causing irreparable damage to the structure and ultimately the function of the cell [29]. One method to prevent the formation of ice crystals and the associated damage is cryopreservation by vitrification [30]. In this process, sperm cryopreservation is performed more quickly and with greater safety than in other methods [31,32]. The cryopreservation process plays a role in the production of ROS by accelerating the conversion of anion superoxide to hydrogen peroxide [33]. Although a

**Table 1.** Effect of *Ceratonia siliqua* concentrations 20, 30, and 40 µg/mL added to cryopreservation/thawing media on motility parameters of human spermatozoa

Groups	Fresh-C	Frozen-C	<i>C. siliqua</i> 20 µg/mL	<i>C. siliqua</i> 30 µg/mL	<i>C. siliqua</i> 40 µg/mL
Motility	45.12 ± 11.10	21.51 ± 8.33 <sup>a)</sup>	31.75 ± 7.13 <sup>b)</sup>	25.30 ± 7.12	25.66 ± 7.91
Progressive motility	40.52 ± 5.21	16.63 ± 5.45 <sup>a)</sup>	41.89 ± 5.14 <sup>b)</sup>	20.55 ± 5.71	21.09 ± 5.41
VCL	63.12 ± 1.16	56.62 ± 0.71 <sup>a)</sup>	59.40 ± 0.79	58.12 ± 0.87	56.33 ± 0.71
VSL	45.91 ± 0.69	37.22 ± 0.51 <sup>a)</sup>	38.17 ± 0.64	36.59 ± 0.58	35.12 ± 0.50
VAP	50.45 ± 0.69	43.64 ± 0.60 <sup>a)</sup>	45.92 ± 0.66	44.12 ± 0.59	43.52 ± 0.67
LIN	59.95 ± 0.62	50.60 ± 0.63 <sup>a)</sup>	51.25 ± 0.57	49.44 ± 0.60	49.75 ± 0.69
SRT	70.30 ± 0.35	55.72 ± 0.31 <sup>a)</sup>	57.11 ± 0.36	54.61 ± 0.30	54.23 ± 0.39
BCF	10.03 ± 0.06	8.71 ± 0.06 <sup>a)</sup>	8.91 ± 0.05	8.77 ± 0.06	8.62 ± 0.06
ALH	9.73 ± 0.05	6.25 ± 0.05 <sup>a)</sup>	6.63 ± 0.06	6.20 ± 0.05	6.22 ± 0.05

Values are presented as mean ± standard error of the mean.

C, control; VCL, curvilinear velocity; VSL, straight-line velocity; VAP, average path velocity; LIN, linearity; SRT, straightness; BCF, beat cross frequency; ALH, amplitude of lateral head displacement.

<sup>a)</sup> $p < 0.05$  significant differences vs. fresh control group; <sup>b)</sup> $p < 0.05$  significant differences vs. frozen control group.

**Table 2.** Effect of *Ceratonia siliqua* concentrations 20, 30, and 40 µg/mL added to cryopreservation/thawing media on ROS intracellular, PMI, viability, MMP, and acrosome reaction of human spermatozoa

Groups	Fresh-C	Frozen-C	<i>C. siliqua</i> 20 µg/mL	<i>C. siliqua</i> 30 µg/mL	<i>C. siliqua</i> 40 µg/mL
ROS intracellular	40.1 ± 6.3	59.1 ± 6.1 <sup>a)</sup>	52.0 ± 6.5 <sup>b)</sup>	52.3 ± 6.1 <sup>b)</sup>	51.3 ± 6.0 <sup>b)</sup>
PMI	53.1 ± 3.5	24.4 ± 3.9 <sup>a)</sup>	55.1 ± 3.4 <sup>b)</sup>	41.3 ± 3.9 <sup>b)</sup>	41.9 ± 3.5 <sup>b)</sup>
Viability	71.3 ± 1.5	45.3 ± 1.09 <sup>a)</sup>	51.6 ± 1.8 <sup>b)</sup>	50.5 ± 1.3 <sup>b)</sup>	50.9 ± 1.3 <sup>b)</sup>
MMP	65.5 ± 4.2	33.6 ± 4.7 <sup>a)</sup>	42.5 ± 4.9 <sup>b)</sup>	44.2 ± 4.9 <sup>b)</sup>	42.9 ± 4.5 <sup>b)</sup>
Acrosome reaction	77.2 ± 5.5	56.5 ± 5.1 <sup>a)</sup>	61.5 ± 5.9 <sup>b)</sup>	60.3 ± 5.2 <sup>b)</sup>	60.9 ± 5.7 <sup>b)</sup>

Values are presented as mean ± standard error of the mean.

ROS, reactive oxygen species; PMI, plasma membrane integrity; MMP, mitochondrial membrane potential; C, control.

<sup>a)</sup> $p < 0.05$  significant differences vs. fresh control group; <sup>b)</sup> $p < 0.05$  significant differences vs. frozen control group.

moderate ROS level is required for optimal sperm function, high levels are associated with the impairment of function [34]. ROS produced as a result of cryopreservation can damage the lipid, protein, and DNA structures of the sperm [35].

Studies have shown that cryopreservation reduces the activity of the sperm antioxidant system, leading to impaired cell motility, integrity, and membrane fluidity [33]. *C. siliqua* extract can protect against oxidative stress [15,36]. Various studies have indicated that the protective effect of this plant extract is due to its strong antioxidant properties [12,19,37]. The results of the current study confirm the protective properties of this extract against oxidative stress, as the intracellular ROS level was significantly reduced at all concentrations of *C. siliqua* extract. Although no previous research had directly examined the intracellular ROS level in sperm treated with *C. siliqua* extract, the improvement of ROS-dependent parameters in other studies aligns with this result. Previous studies have shown that *C. siliqua* extract can increase the activity of antioxidant enzymes such as catalase and superoxide dismutase [15], so the reduction of intracellular ROS during cryopreservation can be explained by the capacity of *C. siliqua* extract to stimulate these enzymes.

Notably, mitochondrial activity in oxidative phosphorylation can be a source of ROS production [38]. At all three concentrations of *C. siliqua* extract, the sperm MMP was significantly increased relative to the cryopreservation control group, most likely by reducing the level of intracellular ROS. Motility is an essential sperm parameter that can have a positive or negative effect on fertility. In the current study, the total and progressive movement in the cryopreservation control group were significantly lower than in the fresh group, likely due to the sensitivity of the adenosine triphosphate-dependent sodium-potassium pump and the consequent leakage of ions related to movement [39]. In the sperm treated with the extract, the total and progressive movement were improved compared to the cryopreservation control group, most notably in the sperm treated with 20 µg/mL of the extract. These findings align with the results of Sabzeie et al. [19] and Faramarzi et al. [12]. Like the present study, those studies revealed not only a positive effect of *C. siliqua* extract on motility, but also an optimal extract concentration of 20 µg/mL. This is likely because *C. siliqua* extract has a greater capacity to reduce superoxide anions at 20 µg/mL than at other concentrations. However, for the other motor characteristics (average path velocity, curvilinear velocity, straight-line velocity, mean linearity, straightness, amplitude of lateral head displacement, and beat cross frequency), we found no significant differences between the concentrations of 20, 30, and 40 µg/mL or in comparison with the cryopreservation control group, which contradicts the study of Sabzeie et al. [19]. Motility is a known indicator of survival and of other health parameters such as PMI, mitochondria health, and even sperm DNA damage.

Our results showed that the presence of *C. siliqua* extract in cryopreservation can help increase sperm viability. This finding supports the results of the Faramarzi et al. [12] study, which revealed a higher percentage of sperm survival in groups treated with the extract than in the control group; additionally, the survival rate of sperm treated with 20 µg/mL was significantly higher than among the control group or the sperm treated with 10 or 40 µg/mL. The improvement of this parameter can likely be attributed to the antioxidant contents of gallic acid, chlorogenic acid, cinnamic acid, and caffeic acid in *C. siliqua* extract. Generally, these compounds have been shown to reduce intracellular ROS and increase sperm viability. In the current study, the results of the HOS test indicated that the greatest membrane integrity was present in the group treated with 20 µg/mL of extract, while this parameter was not significantly different in the groups with concentrations of 30 and 40 µg/mL. This finding is completely consistent with the result of Sabzeie et al. [19]. In our study, the presence of *C. siliqua* extract at all concentrations (20, 30, and 40 µg/mL) led to a significant increase in the acrosome reaction compared to the cryopreservation control group. This result can be attributed to the antioxidant roles of this extract in countering the damage caused by oxidative stress as well as in osmotic balance regulation, membrane preservation, conjugation, and calcium regulation. Accordingly, the role of extracellular antioxidants in improving sperm membrane resistance can be explained by the reduction of lipid peroxidation [40].

In conclusion, *C. siliqua* extract supplements at concentrations of 20, 30, and 40 µg/mL improve sperm motility, viability, cell membrane and mitochondrial potential, intracellular ROS, and the acrosome reaction. A concentration of 20 µg/mL has a particularly high capacity to reduce intracellular ROS, oxidative stress, and its products, eventually improving ROS-affected parameters such as motility, PMI, MMP, and the acrosome reaction.

## Conflict of interest

No potential conflict of interest relevant to this article was reported.

## ORCID

Tooba Farazmand

<https://orcid.org/0000-0002-4380-010x>

Erfan Shahandeh

<https://orcid.org/0000-0002-7004-4514>

## Author contributions

Conceptualization: YK, ES. Data curation: FM. Formal analysis: TF. Funding acquisition: ES. Methodology: YK. Project administration: TF.

Visualization: TF. Writing-original draft: YK. Writing-review & editing: FM, YK.

## References

- Oehninger S, Duru NK, Srisombut C, Morshedi M. Assessment of sperm cryodamage and strategies to improve outcome. *Mol Cell Endocrinol* 2000;169:3–10.
- Braga DP, Setti AS, Figueira RC, Iaconelli A Jr, Borges E Jr. The negative influence of sperm cryopreservation on the quality and development of the embryo depends on the morphology of the oocyte. *Andrology* 2015;3:723–8.
- Shufaro Y, Schenker JG. Cryopreservation of human genetic material. *Ann N Y Acad Sci* 2010;1205:220–4.
- Tamburrino L, Cambi M, Marchiani S, Manigrasso I, Degl'Innocenti S, Forti G, et al. Sperm DNA fragmentation in cryopreserved samples from subjects with different cancers. *Reprod Fertil Dev* 2017;29:637–45.
- Di Santo M, Tarozzi N, Nadalini M, Borini A. Human sperm cryopreservation: update on techniques, effect on DNA integrity, and implications for ART. *Adv Urol* 2012;2012:854837.
- Gupta S, Agarwal A, Sharma R, Ahmady A. Recovery, preparation, storage and utilization of spermatozoa for fertility preservation in cancer patients and sub-fertile men. *J Reprod Stem Cell Biotechnol* 2010;1:150–68.
- Amidi F, Pazhohan A, Shabani Nashtaei M, Khodarahmian M, Nekoonam S. The role of antioxidants in sperm freezing: a review. *Cell Tissue Bank* 2016;17:745–56.
- Kopeika J, Zhang T, Rawson DM, Elgar G. Effect of cryopreservation on mitochondrial DNA of zebrafish (*Danio rerio*) blastomere cells. *Mutat Res* 2005;570:49–61.
- Layek SS, Mohanty TK, Kumaresan A, Parks JE. Cryopreservation of bull semen: evolution from egg yolk based to soybean based extenders. *Anim Reprod Sci* 2016;172:1–9.
- Hezavehei M, Sharafi M, Kouchesfahani HM, Henkel R, Agarwal A, Esmaeili V, et al. Sperm cryopreservation: a review on current molecular cryobiology and advanced approaches. *Reprod Biomed Online* 2018;37:327–39.
- Mohammadi F, Varanloo N, Heydari Nasrabadi M, Vatannejad A, Amjadi FS, Javedani Masroor M, et al. Supplementation of sperm freezing medium with myoinositol improve human sperm parameters and protects it against DNA fragmentation and apoptosis. *Cell Tissue Bank* 2019;20:77–86.
- Faramarzi A, Aghaz F, Golestan Jahromi M, Bakhtiari M, Khazaei M. Does supplementation of sperm freezing/thawing media with *Ceratonia siliqua* improve detrimental effect of cryopreservation on sperm parameters and chromatin quality in normozoospermic specimens? *Cell Tissue Bank* 2019;20:403–9.
- Forouzanfar M, Sharafi M, Hosseini SM, Ostadhosseini S, Hajian M, Hosseini L, et al. In vitro comparison of egg yolk-based and soybean lecithin-based extenders for cryopreservation of ram semen. *Theriogenology* 2010;73:480–7.
- Faramarzi A, Aghaz F, Bakhtiari M, Roshankhah S, Rashidi Z, Khazaei M. *Ceratonia siliqua* (Carob) extract improved in vitro development of vitrified-warmed mouse germinal vesicle oocytes: assessment of possible mechanism. *Cell Tissue Bank* 2021;22:137–44.
- Vafaei A, Mohammadi S, Fazel A, Soukhtanloo M, Mohammadi-pour A, Beheshti F. Effects of carob (*Ceratonia siliqua*) on sperm quality, testicular structure, testosterone level and oxidative stress in busulfan-induced infertile mice. *Pharma Sci* 2018;24:104–11.
- Pan B, Yang H, Wu Z, Qazi IH, Liu G, Han H, et al. Melatonin improves parthenogenetic development of vitrified-warmed mouse oocytes potentially by promoting G1/S cell cycle progression. *Int J Mol Sci* 2018;19:4029.
- Khazaei M, Pazhouhi M, Khazaei S. Evaluation of hydro-alcoholic extract of *Trifolium pratense* L. for its anti-cancer potential on U87MG cell line. *Cell J* 2018;20:412–21.
- Sadeghzadeh F, Sadeghzadeh A, Changizi-Ashtiyani S, Bakhshi S, Mashayekhi FJ, Mashayekhi M, et al. The effect of hydro-alcoholic extract of *Ceratonia Siliqua* L. on spermatogenesis index in rats treated with cyclophosphamide: an experimental study. *Int J Reprod Biomed* 2020;18:295–306.
- Sabzeie MM, Soleimanzadeh A, Ayen E. The effect of *Ceratonia siliqua* aqueous extract on canine semen quality on during storage at Refrigerator. *Vet Res Biol Prod* 2021;34:120–10.
- Ghaleno LR, Valojerdi MR, Janzamin E, Chehrizi M, Sharbatoghli M, Yazdi RS. Evaluation of conventional semen parameters, intracellular reactive oxygen species, DNA fragmentation and dysfunction of mitochondrial membrane potential after semen preparation techniques: a flow cytometric study. *Arch Gynecol Obstet* 2014;289:173–80.
- Taher-Mofrad SM, Topraggaleh TR, Ziarati N, Bucak MN, Nouri M, Seifi S, et al. Knockout serum replacement is an efficient serum substitute for cryopreservation of human spermatozoa. *Cryobiology* 2020;92:208–14.
- Aitken RJ, Clarkson JS. Significance of reactive oxygen species and antioxidants in defining the efficacy of sperm preparation techniques. *J Androl* 1988;9:367–76.
- Seify M, Zarabadipour M, Ghaleno LR, Alizadeh A, Rezazadeh Valojerdi M. The anti-oxidant roles of taurine and hypotaurine on acrosome integrity, HBA and HSPA2 of the human sperm during vitrification and post warming in two different temperature. *Cryobiology* 2019;90:89–95.

24. Ziarati N, Topraggaleh TR, Rahimizadeh P, Montazeri L, Maroufzadeh S, Sadighi Gilani MA, et al. Micro-quantity straw as a carrier for cryopreservation of oligozoospermic semen samples: effects of storage times and cryoprotectant. *Cryobiology* 2019;86:65–70.
25. Bucak MN, Bodu M, Baspinar N, Gungor S, Ili P, Acibaeva B, et al. Influence of ellagic acid and ebselen on sperm and oxidative stress parameters during liquid preservation of ram semen. *Cell J* 2019;21:7–13.
26. Koohestanidehaghi Y, Torkamanpari M, Shirmohamadi Z, Lorian K, Vatankhah M. The effect of cysteine and glutamine on human sperm functional parameters during vitrification. *Andrologia* 2021;53:e13870.
27. Torkamanpari M, Ghorbani F, Lorian K, Koohestanidehaghi Y. The effects of purslane (*Portulaca oleracea*) and fennel (*Foeniculum vulgare* Mill) hydroalcoholic extracts on the functional parameters of human spermatozoa after vitrification. *Clin Exp Reprod Med* 2023;50:78–85.
28. Nasiri Z, Ghorbani F, Seify M, Sharbati A. Effect of aqueous *Nigella sativa* extract on the functional parameters of post-thaw human spermatozoa during vitrification. *Clin Exp Reprod Med* 2022;49:110–6.
29. Tiwari A, Tekcan M, Sati L, Murk W, Stronk J, Huszar G. A new media without animal component for sperm cryopreservation: motility and various attributes affecting paternal contribution of sperm. *J Assist Reprod Genet* 2017;34:647–57.
30. Arav A, Natan Y, Kalo D, Komsky-Elbaz A, Roth Z, Levi-Setti PE, et al. A new, simple, automatic vitrification device: preliminary results with murine and bovine oocytes and embryos. *J Assist Reprod Genet* 2018;35:1161–8.
31. Horta F, Alzobi H, Jitanantawittaya S, Catt S, Chen P, Pangestu M, et al. Minimal volume vitrification of epididymal spermatozoa results in successful in vitro fertilization and embryo development in mice. *Asian J Androl* 2017;19:107–12.
32. Spis E, Bushkovskaia A, Isachenko E, Todorov P, Sanchez R, Skopets V, et al. Conventional freezing vs. cryoprotectant-free vitrification of epididymal (MESA) and testicular (TESE) spermatozoa: three live births. *Cryobiology* 2019;90:100–2.
33. Ghorbani F, Nasiri Z, Koohestanidehaghi Y, Lorian K. The antioxidant roles of L-carnitine and N-acetyl cysteine against oxidative stress on human sperm functional parameters during vitrification. *Clin Exp Reprod Med* 2021;48:316–21.
34. Lone SA, Prasad JK, Ghosh SK, Das GK, Balamurugan B, Verma MR. Study on correlation of sperm quality parameters with antioxidant and oxidant status of buffalo bull semen during various stages of cryopreservation. *Andrologia* 2018;50:e12970.
35. Bustamante Filho IC, Pederzoli CD, Sgaravatti AM, Gregory RM, Dutra Filho CS, Jobim MI, et al. Skim milk-egg yolk based semen extender compensates for non-enzymatic antioxidant activity loss during equine semen cryopreservation. *Anim Reprod* 2018;6:392–9.
36. Rtibi K, Selmi S, Grami D, Saidani K, Sebai H, Amri M, et al. *Ceratonia siliqua* L. (immature carob bean) inhibits intestinal glucose absorption, improves glucose tolerance and protects against alloxan-induced diabetes in rat. *J Sci Food Agric* 2017;97:2664–70.
37. Faramarzi A, Aghaz F, Bakhtiari M, Khazaei M. In vitro application of *Ceratonia siliqua* improved sperm parameters and chromatin quality after vitrification in normozoospermic aged men. *Middle East Fertil Soc J* 2020;24:6.
38. He B, Guo H, Gong Y, Zhao R. Lipopolysaccharide-induced mitochondrial dysfunction in boar sperm is mediated by activation of oxidative phosphorylation. *Theriogenology* 2017;87:1–8.
39. Saito K, Kinoshita Y, Kanno H, Iwasaki A. The role of potassium ion and extracellular alkalization in reinitiation of human spermatozoa preserved in electrolyte-free solution at 4 degrees C. *Fertil Steril* 1996;65:1214–8.
40. Dawson R Jr, Biasetti M, Messina S, Dominy J. The cytoprotective role of taurine in exercise-induced muscle injury. *Amino Acids* 2002;22:309–24.