

Better isolation, proliferation and differentiation of human adipose-derived mesenchymal stem cells using human serum

Atena Sadat Ghoreishi (1), Ehsan Iranmanesh (2), Hosna Rastegarpouyani (3), Sogand Mokhtarian (4), Zahra Poshtchaman (5), Zeinab Sadat Javadi (6), Alireza Khoshdel (7,8)

(1) Department of Clinical Biochemistry, Faculty of Para-Medicine, Jiroft University of Medical Sciences, Jiroft, Iran; (2) Neuroscience Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran; (3) Department of Biological Science, Florida State University, Tallahassee, FL, USA Institute for Molecular Biophysics, Florida State University, Tallahassee, FL, USA; (4) Department of Cell and Molecular Biology, Islamic Azad University of Shahr-e-Qods, Tehran, Iran; (5) MSc of Critical care Nursing, Department of Nursing, Esfarayen Faculty of Medical Sciences, Esfarayen, Iran; (6) Yazd hospital Mehrab Shohada, Shahid Sadoughi University of Medical Sciences, Yazd, Iran; (7) Department of Clinical Biochemistry, Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran; (8) Nervous System Stem Cells Research Center, Semnan University of Medical Sciences, Semnan, Iran.

This article is distributed under the terms of the Creative Commons Attribution Noncommercial License (CC BY-NC 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Abstract

Mesenchymal stem cells have many applications in medicine. Attention to the proliferation and differentiation of stem cell differentiation is an important issue. The aim of this study was to investigate the possibility of optimal isolation, proliferation, and differentiation of adipose tissue-derived mesenchymal stem cells (ADSCs) using human serum. Human serum (HS) was obtained from the venous blood of eight healthy individuals. The rate of proliferation and differentiation of ADSCs and expression of surface markers was assessed by flow cytometry. Bone differentiation was assessed using Alizarin Red staining. Data were analyzed using statistical software. Over time, HS showed more proliferation than fetal bovine serum (FBS) -enriched cells ($p < 0.05$). Differentiation of ADSCs cells in HS-enriched medium is faster and more pronounced than differentiation in the control group. The expression of surface markers in the medium containing HS was the same as the medium containing FBS where the expression levels of CD105 and CD95 were found to be positive and the expression of CD34 and CD45 was negative. Due to the better proliferation of adipose tissue-derived mesenchymal cells in the medium containing HS than FBS, it is suggested that human serum be used in future clinical studies. Also, HS is healthier, safer, more accessible, and more affordable than FBS.

Key Words: Replication; differentiation; mesenchymal stem cells; human serum; human adipose-derived stem cells (ADSCs).

Eur J Transl Myol 33 (1) 10834, 2023 doi: 10.4081/ejtm.2023.18340

So far, many research studies have been conducted in the field of medicine, most of which are in the field of common and infectious diseases. For instance, various studies have been conducted in the field of cancer,¹⁻⁵ and in recent years, with the outbreak of coronavirus,⁶⁻⁸ studies have mainly gone in this direction. However, the treatment of these diseases has always been very significant. Additionally, much research has been conducted regarding medicine and health statistics to prove the importance of the healthcare system.⁹⁻¹² One of

these methods is cell therapy using stem cells and extracellular vesicles derived from these cells. Stem cells have many applications in medicine.^{13,14} Mesenchymal stem cells are commercially used for three reasons: first, they can be located in areas of inflammation and tissue damage, such as wounds or tumors. Second, mesenchymal stem cells have immunosuppressive properties and can modulate innate and adaptive immune responses. Third, these cells have the ability to support other cells with paracrine factors.¹³⁻¹⁵ Among these, mesenchymal stem cells are of great interest.

Human mesenchymal stem cells and human serum

Eur J Transl Myol 33 (1) 10834, 2023 doi: 10.4081/ejtm.2023.10834

Mesenchymal stem cells have the power to differentiate into different cells and are also able to become undifferentiated cells similar to their own. These cells have the ability to differentiate into connective tissue lineages, including adipocytes, chondrocytes, and osteocytes. Mesenchymal stem cells are capable of regenerating their divisions for a long time and at the same time maintaining their differentiation power.^{16,17} The clinical application of stem cells in the treatment of brain injury, Parkinson's, myocardial infarction, incomplete osteogenesis, fracture healing, and tendon rupture, as well as treatments of joint, liver and other disorders have been well demonstrated.^{18,19}

One of important sources of potent mesenchymal stem cells is adipose tissue, which is considered a very suitable source of stem cells.^{20,21} Adipose tissue-derived stem cells (ADSCs) are easily harvested and multiplied, are effective and non-invasive, and have a high capacity for multiplication and proliferation. These types of autologous stem cells significantly reduce donation complications.^{22,23}

On the other hand, literature findings suggest that use of animal serum in clinical cases of cell therapy is not a completely safe choice. Indeed, animal serum may cause immune reactions and infections.²⁴⁻²⁶ Providing the safest and most cost-effective serum with optimal properties and greater safety for cell therapy is an important issue.²⁷ Human serum (HS) seems to be a suitable alternative, because of its high availability, safety and rapid collection. Human serum contains many useful proteins, including growth factors and immunoglobulins, inducing a wide variety of positive effects on cell culture.²⁸

Studies on ADSC have found that HS can be used to produce and proliferate fat stem cells, reporting that human ADSC proliferation in growth medium containing HS is higher than in culture medium containing fetal bovine serum (FBS). In fact, proteins and other nutrient factors in human serum affect the expression of inflammatory agents in stem cells.³⁰⁻³²

Quantitative studies have been performed on various aspects of the advantage of human serum over animal serum, so the aim of this study was to confirm or not the effects of human serum for better isolation, proliferation and differentiation of ADSC in cell culture medium.

Materials and Methods

Preparation of medium and HS serum

A total of 100 cc of venous blood was taken from 8 healthy adult donors with written consent. The prepared samples were centrifuged at 3000 rpm for 8 minutes and the serum was separated. The isolated serum was incubated for 56 minutes at 56 ° C to inactivate. It was then placed at -80 ° C for later use.

Cell extraction and culture

ADSCs were prepared from liposuctioned human adipose tissue. 60 g of adipose tissue sample was transferred to the culture room in a sterile container. In order to remove blood cells, fat samples were washed several times with Phosphate-Buffered Saline (PBS) solution (Shellmax) and penicillin and streptomycin (Gibco) antibiotics were added. The adipose tissue was then divided into small pieces and mixed with 3 ml of type I collagenase enzyme (0.075%) (Gibco) dissolved in PBS. After digestion, the small pieces were placed in a




Fig 1. Difference in the number of cells on consecutive days of culture in both mediums enriched with HS and FBS

37-degree incubator for 40 minutes, during which time the sample was shaken every 8 minutes for further digestion of the tissue.

The sample was then divided into two parts, one was added to DMEM (Gibco) medium enriched with 10% HS (human serum) and the other to DMEM (Gibco) medium enriched with 10% FBS (animal serum). The cells were then centrifuged at 1200 rpm for 6 minutes. After the addition of penicillin and streptomycin antibiotics to the solution containing adipose tissue mesenchymal cells. Finally, cultures containing HS and FBS were incubated at 37 ° C with 5% CO₂ and 90% humidity.

It is noteworthy that adipose tissue was collected after receiving written consent from the volunteer of lipolysis and this operation was performed by a surgeon in the operating room under completely sterile conditions.

Cell proliferation of ADSCs

The culture medium of ADSCs containing HS or FBS was replaced with a fresh medium every three days after incubation with the mentioned conditions. After cell density reached 90-70% by trypsinization (trypsin / EDTA; Sigma) of the culture plate, the isolated cells were used for cell passage or subsequent analysis.³³

Stem cell proliferation rates in both samples containing HS or FBS were assessed after the second passage. Some cells with a density of 103 were selected and cultured in

4-well cell culture plates according to the conditions of the cultured cells. Cells were examined for proliferation on days 3, 5, 7, and 9. Cell passage

As the cells grew in the culture flask, the supernatant was discarded and the cells were washed with PBS. The cells were then added to the trypsin flask and incubated in a 37 ° C incubator for 3 minutes. After separating the cells from the flask, DMEM containing HS or FBS was added with trypsin to inactivate them. The solution containing the cells was then centrifuged at 1000 rpm for 6 minutes. In order to count the cells, 10 ml of homogeneous medium was cultured in 4-well cell culture plates 3 times each time in each cell passage.

In each passage, their number was counted using a *Neobar* lam and an electronic reed counter (Japan, Cis Max) in the capillary position.

Differentiation of adipocytes and cellular osteocytes

To evaluate the ability to differentiate ADSC in HS in the adipogenic differentiation, cells were incubated in DMEM supplemented with dexamethasone M μ 1, insulin M μ 10, indomethacin M μ 200, and isobutyrate methylxanthine M μ 0.5 in either medium supplemented with HS or FBS. About two weeks later, ADSC was fixed in 10% formalin neutral buffer for 25 minutes and stained with Oil red O. The dye reaction was examined under an inverted microscope.




Fig 2. Specific staining of cultured cells containing medium enriched with HS and FBS. A. Control group (FBS), b) The first group of human serum (day 9), c) The second group of human serum (day 12), d) The third group of human serum (15th day), (10x magnification).

To differentiate the bone class, ADSC was cultured on culture plates. First, the primary culture medium was discarded. Cells were then incubated in DMEM differential medium containing 7-10 M dexamethasone, 0.2 M ascorbic acid, and 10 mM beta-glycerol phosphate in either medium supplemented with HS or FBS for 21 days. The existing culture medium was changed based on ocular vision or using a microscope every 3 days. Differentiation was assessed by Alizarin Red staining to detect calcium stores.

Data analysis

Frequency and percentage were used to express qualitative variables. T-test was used to evaluate the variables. SPSS software version 25 was used for analysis. A p value less than 0.05 was considered to be statistically significant.

Results

Cell proliferation of ADSCs

Cell proliferation findings showed that ADSCs in HS-enriched medium in the early days were not significantly different from FBS-enriched cells. However, serum containing HS showed significantly more proliferation over time than FBS-enriched cells ($p < 0.05$; Figure 1). The difference between the seventh and ninth days was statistically significant, where the groups had a significant difference on different days ($p < 0.05$).

Ability to differentiate into cellular osteocytes

From the tenth day, calcium deposition, which was a characteristic of bone differentiation, was observed in the culture medium and the differentiation reached the desired level on the 15th day. According to Figure 2, it was observed that the differentiation of ADSCs cells in the medium enriched with HS is faster and more pronounced than the differentiation in the control group.

Expression of cell surface markers

In flow cytometric analysis performed on cells harvested from the fourth passage, the expression pattern of surface

markers in proliferating cells in media enriched with HS and FBS was observed. The expression of surface markers in the medium containing HS was the same as the medium containing FBS where the expression levels of CD105 and CD95 were seen and the expression of CD 34 and CD45 was negative (Figure 2).

Ability to differentiate cellular adipocytes

Cells cultured with HS and FBS-enriched media that were affected by adipocyte differentiation had changed from a spindle to a spherical shape in both groups and contained fat vacuoles (Figure 3).

Discussion

Much research has been done on the differentiation of mesenchymal stem cells. However, limited studies have been performed so far on the use of HS for better proliferation of ADSCs. The present study seeks to optimize the proliferation and differentiation of mesenchymal stem cells using human serum. The findings of the present study showed that the proliferation of ADSCs in the medium containing HS is better and more efficient than in the medium containing FBS. Studies by Lindroos et al.²⁹ and Bieback et al. also stated that the optimal percentage of human ADSC proliferation in a human serum-containing growth medium was higher than that of an FBS culture medium.³⁰

Josh et al. showed that human ADSC growth in HS-supplemented medium was faster than that of FBS-supplemented medium.³³ Another study found that the use of self-derived serum was less dangerous than the use of animal-derived serum and reduced the harms of using FBS for ADSC proliferation.³⁴

In a study by Bahn et al.,³⁵ it was reported that ADSC cells proliferated and grew faster in culture medium containing human serum than in culture medium containing FBS. Thus, human serum can potentially be used instead of FBS in ADSC medium. de Paula et al. (2013)³⁶ stated that human serum is a promising supplement for the proliferation of ADSC cells in the laboratory.




Fig 3. Specific oil red O staining on cultured cells containing media enriched with HS and FBS. A) Control group (FBS), b) Human serum group.

A study by Koellensperger et al.³⁷ showed that culturing ADSC cells in human serum is a viable alternative to FBS. In another study in 2021, the effect of FBS, human serum and human platelets on mesenchymal stem cells was investigated by examining the behavior of mesenchymal stem cells (metabolic activity, proliferation, morphology, and cytokine production). The results of this study showed that human serum (autologous and allogeneic) and released platelet lysates are promising and relatively good alternatives for MSC culture.³⁸ This study confirms that differentiation of ADSCs cells in HS-enriched medium was faster and more pronounced than the differentiation in the control group. Studies by Dreher et al. (2013)³⁹ showed that the more differentiating effects of HS on ADSC cell culture were greater than those of FBS. In general, similar studies have focused little on this issue. Given the role of differentiation in stem cell function, this is a very important issue.

Of course, one of the limitations that has been raised in various studies related to human serum is the condition of donors in terms of age, immune system, and serum quality. Some studies have suggested that serum donor conditions affect mesenchymal cell metabolism. Some factors including different serum quality (the content of growth factors and immune elements), serum obtained from adult donors (exposure of the immune system to various stimuli throughout life), and serum extracted from younger people (more naive immune system) have been mentioned to be capable of affecting the mesenchymal cell metabolism.^{40,41}

To overcome this limitations, in the present study, the serum of 8 people was used and donors were selected from healthy adults.

In conclusion, our study confirm that mesenchymal cell proliferation and bone differentiation were higher in HS medium than in FBS. Due to better proliferation of adipose tissue-derived mesenchymal cells in the medium containing HS than FBS, it is suggested that human serum can be used in future clinical studies. Furthermore, study of compounds in human serum can be effectively used for better optimization of culture media.

List of acronyms

ADSCs - adipose tissue-derived mesenchymal stem cells

CD34 - Cluster of Differentiation 34

CD45 - Cluster of Differentiation 45

CD105 - Cluster of Differentiation 105

FBS - fetal bovine serum

HS - Human serum

Contributions of Authors

Acquisition of main funding: Rafsanjan University of Medical Sciences.

Acknowledgments

We apologize to colleagues whose research papers have not been cited due to space constraints.

Funding

This work was funded by Rafsanjan University of Medical Sciences.

Conflict of Interest

The authors declare no conflict of interest.

Ethical Publication Statement

I confirm that I have read the Journal's position on ethical publication issues and affirm that this report is consistent with those guidelines.

Corresponding Author

Alireza Khoshdel, Department of Clinical Biochemistry, Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran. Nervous System Stem Cells Research Center. Semnan University of Medical Sciences, Semnan, Iran.

ORCID ID: 0000-0002-2148-4951

E-mail: Alireza.khoshdel@gmail.com

E-mails and ORCID iD of co-authors

Atena Sadat Ghoreishi: atooo4929@Yahoo.com

ORCID iD: 0000-0001-9312-611X

Ehsan Iranmanesh: E_iranmanesh@kmu.ac.ir

ORCID iD: 0000-0003-1342-8551

Hosna Rastegarpouyani: hr18d@fsu.edu

ORCID iD: 0000-0003-1358-9910

Sogand Mokhtarian: sogandmokhtarian1@gmail.com

ORCID iD: 0000-0001-6806-6629

Zahra Poshtchaman: zahraposhtchaman@yahoo.com

ORCID iD: 0000-0002-3712-1493

Zeinab Sadat Javadi: javadi.ieshtiri9@gmail.com

ORCID iD: 0000-0001-7932-0135

References

1. Ebrahimi N, Adelian S, Shakerian S, Afshinpour M, Chaleshtori SR, Rostami N, Rezaei-Tazangi F, Beiranvand S, Hamblin MR, Aref AR. Crosstalk between ferroptosis and the epithelial-mesenchymal transition: Implications for inflammation and cancer therapy. *Cytokine Growth Factor Rev.* 2022 Apr;64:33-45. doi: 10.1016/j.cytogfr.2022.01.006.
2. Kazemi M, Emami J, Hasanazadeh F, Minaiyan M, Mirian M, Lavasanifar A, Mokhtari M. *In Vitro* and *In Vivo* Evaluation of Novel DTX-Loaded Multifunctional Heparin-Based Polymeric Micelles Targeting Folate Receptors and Endosomes. *Recent Pat Anticancer Drug Discov.* 2020;15(4):341-359. doi: 10.2174/1574892815666201006124604.
3. Kazemi, M., Emami, J., Hasanazadeh, F., Minaiyan, M., Mirian, M., & Lavasanifar, A. Pegylated multifunctional pH-responsive targeted polymeric micelles for ovarian cancer therapy: synthesis, characterization and pharmacokinetic study. *International Journal of Polymeric Materials and*

- Polymeric Biomaterials. 2021 Sep;70(14):1012-1026. Doi:10.1080/00914037.2020.1776282.
4. Molani, S., Madadi, M., & Wilkes, W. A partially observable Markov chain framework to estimate overdiagnosis risk in breast cancer screening: Incorporating uncertainty in patients adherence behaviors. *Omega*, Elsevier. 2019 Dec;89(C),40-53. doi: 10.1016/j.omega.2018.09.009.
 5. Molani, S., Madadi, M., & Williams, D. Investigating the effectiveness of supplemental breast cancer screening tests considering radiologists' bias. medRxiv. 2020.12.16.20248373. doi:10.1101/2020.12.16.20248373
 6. Khorsandi Z, Afshinpour M, Molaei F, Askandar RH, Keshavarzipour F, Abbasi M, Sadeghi-Aliabadi H. Design and synthesis of novel phe-phe hydroxyethylene derivatives as potential coronavirus main protease inhibitors. *J Biomol Struct Dyn*. 2022 Oct;40(17):7940-7948. doi: 10.1080/07391102.2021.1905549.
 7. Norouzi M, Hashemi M, Pouri Z. The Question of Global Society in Post-Corona Time: Towards a Paradigm Shift. *Int J Community Wellbeing*. 2021;4(3):339-343. doi: 10.1007/s42413-021-00141-7.
 8. Khorani M, Bobe G, Matthews DG, Magana AA, Caruso M, Gray NE, Quinn JF, Stevens JF, Soumyanath A, Maier CS. The Impact of the hAPP695SW Transgene and Associated Amyloid- β Accumulation on Murine Hippocampal Biochemical Pathways. *J Alzheimers Dis*. 2022;85(4):1601-1619. doi: 10.3233/JAD-215084.
 9. Mostoufi A, Chamkouri N, Kordrostami S, Alghasibabaahmadi E, Mojaddami A. 3-Hydroxypyrimidine-2, 4-dione Derivatives as HIV Reverse Transcriptase-Associated RNase H Inhibitors: QSAR Analysis and Molecular Docking Studies. *Iran J Pharm Res*. 2020 Winter;19(1):84-97. doi: 10.22037/ijpr.2020.1101004.
 10. Tomar N, Sadri S, Cowley AW Jr, Yang C, Quryshi N, Pannala VR, Audi SH, Dash RK. A thermodynamically-constrained mathematical model for the kinetics and regulation of NADPH oxidase 2 complex-mediated electron transfer and superoxide production. *Free Radic Biol Med*. 2019 Apr;134:581-597. doi: 10.1016/j.freeradbiomed.2019.02.003.
 11. Rafiepour, H., Abdollah Zadeh, A., Moradan, A., Salekshahrezaee, Z. Study of Genes Associated With Parkinson Disease Using Feature Selection. *Journal of Bioengineering Research*. 2020 Dec;2(4),1-11. Doi:10.22034/JBR.2020.251812.1035.
 12. Rezapour-Nasrabad, R. Transitional care model: managing the experience of hospital at home. *Electronic Journal of General Medicine* 2018 Aug;15(5):em73. Doi:10.29333/ejgm/93445
 13. Foo JB, Looi QH, Chong PP, Hassan NH, Yeo GEC, Ng CY, Koh B, How CW, Lee SH, Law JX. Comparing the Therapeutic Potential of Stem Cells and their Secretory Products in Regenerative Medicine. *Stem Cells Int*. 2021 Aug 19;2021:2616807. doi: 10.1155/2021/2616807.
 14. Abdelgawad M, Allam S, Abdelmonaem Shaheen M, Ali Hussein M, Azmy Elkot H, Gaber A, et al. An overview of COVID-19 treatment: possible candidates based on drug repurposing and molecular docking. *Canadian Journal of Medical Sciences*. 2021 May;3(1):10–35. Doi: 10.33844/cjm.2021.60499.
 15. Khosrotehrani, K. Mesenchymal stem cell therapy in skin: why and what for?. *Experimental Dermatology*. 2013 May;22(5):307-10. doi: 10.1111/exd.12141.
 16. Razeghian-Jahromi I, Matta AG, Canitrot R, Zibaenezhad MJ, Razmkhah M, Safari A, Nader V, Roncalli J. Surfing the clinical trials of mesenchymal stem cell therapy in ischemic cardiomyopathy. *Stem Cell Res Ther*. 2021 Jun 23;12(1):361. doi: 10.1186/s13287-021-02443-1.
 17. Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, Moorman MA, Simonetti DW, Craig S, Marshak DR. Multilineage potential of adult human mesenchymal stem cells. *Science*. 1999 Apr 2;284(5411):143-7. doi: 10.1126/science.284.5411.143.
 18. Hung SC, Chen NJ, Hsieh SL, Li H, Ma HL, Lo WH. Isolation and characterization of size-sieved stem cells from human bone marrow. *Stem Cells*. 2002;20(3):249-58. doi: 10.1634/stemcells.20-3-249.
 19. Xie B, Chen S, Xu Y, Han W, Hu R, Chen M, He R, Ding S. Clinical Efficacy and Safety of Human Mesenchymal Stem Cell Therapy for Degenerative Disc Disease: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Stem Cells Int*. 2021 Sep 13;2021:9149315. doi: 10.1155/2021/9149315.
 20. Rodríguez LV, Alfonso Z, Zhang R, Leung J, Wu B, Ignarro LJ. Clonogenic multipotent stem cells in human adipose tissue differentiate into functional smooth muscle cells. *Proc Natl Acad Sci U S A*. 2006 Aug 8;103(32):12167-72. doi: 10.1073/pnas.0604850103.
 21. Bunnell BA, Flaatt M, Gagliardi C, Patel B, Ripoll C. Adipose-derived stem cells: isolation, expansion and differentiation. *Methods*. 2008 Jun;45(2):115-20. doi: 10.1016/j.ymeth.2008.03.006.
 22. Rodríguez LV, Alfonso Z, Zhang R, Leung J, Wu B, Ignarro LJ. Clonogenic multipotent stem cells in human adipose tissue differentiate into functional smooth muscle cells. *Proc Natl Acad Sci U S A*. 2006 Aug 8;103(32):12167-72. doi: 10.1073/pnas.0604850103.

23. Ogawa R. The importance of adipose-derived stem cells and vascularized tissue regeneration in the field of tissue transplantation. *Curr Stem Cell Res Ther.* 2006 Jan;1(1):13-20. doi: 10.2174/157488806775269043.
24. Gstraunthaler, G. Alternatives to the use of fetal bovine serum: serum-free cell culture. *ALTEX.* 2003;20(4):275-81. PMID: 14671707.
25. Martin MJ, Muotri A, Gage F, Varki A. Human embryonic stem cells express an immunogenic nonhuman sialic acid. *Nat Med.* 2005 Feb;11(2):228-32. doi: 10.1038/nm1181.
26. Komoda H, Okura H, Lee CM, Sougawa N, Iwayama T, Hashikawa T, Saga A, Yamamoto-Kakuta A, Ichinose A, Murakami S, Sawa Y, Matsuyama A. Reduction of N-glycolylneuraminic acid xenoantigen on human adipose tissue-derived stromal cells/mesenchymal stem cells leads to safer and more useful cell sources for various stem cell therapies. *Tissue Eng Part A.* 2010 Apr;16(4):1143-55. doi: 10.1089/ten.TEA.2009.0386.
27. Kobayashi T, Watanabe H, Yanagawa T, Tsutsumi S, Kayakabe M, Shinozaki T, Higuchi H, Takagishi K. Motility and growth of human bone-marrow mesenchymal stem cells during ex vivo expansion in autologous serum. *J Bone Joint Surg Br.* 2005 Oct;87(10):1426-33. doi: 10.1302/0301-620X.87B10.16160.
28. Blande IS, Bassaneze V, Lavini-Ramos C, Fae KC, Kalil J, Miyakawa AA, Schettert IT, Krieger JE. Adipose tissue mesenchymal stem cell expansion in animal serum-free medium supplemented with autologous human platelet lysate. *Transfusion.* 2009 Dec;49(12):2680-5. doi: 10.1111/j.1537-2995.2009.02346.x.
29. Lindroos B, Aho KL, Kuokkanen H, Rätty S, Huhtala H, Lemponen R, Yli-Harja O, Suuronen R, Miettinen S. Differential gene expression in adipose stem cells cultured in allogeneic human serum versus fetal bovine serum. *Tissue Eng Part A.* 2010 Jul;16(7):2281-94. doi: 10.1089/ten.TEA.2009.0621.
30. Bieback K, Ha VA, Hecker A, Grassl M, Kinzebach S, Solz H, Sticht C, Klüter H, Bugert P. Altered gene expression in human adipose stem cells cultured with fetal bovine serum compared to human supplements. *Tissue Eng Part A.* 2010 Nov;16(11):3467-84. doi: 10.1089/ten.TEA.2009.0727.
31. Jenhani F, Durand V, Ben Azouna N, Thallet S, Ben Othmen T, Bejaoui M, Domenech J. Human cytokine expression profile in various conditioned media for in vitro expansion bone marrow and umbilical cord blood immunophenotyped mesenchymal stem cells. *Transplant Proc.* 2011 Mar;43(2):639-43. doi: 10.1016/j.transproceed.2011.01.021.
32. Mimura S, Kimura N, Hirata M, Tateyama D, Hayashida M, Umezawa A, Kohara A, Nikawa H, Okamoto T, Furue MK. Growth factor-defined culture medium for human mesenchymal stem cells. *Int J Dev Biol.* 2011;55(2):181-7. doi: 10.1387/ijdb.103232sm.
33. Josh F, Kobe K, Tobita M, Tanaka R, Suzuki K, Ono K, Hyakusoku H, Mizuno H. Accelerated and safe proliferation of human adipose-derived stem cells in medium supplemented with human serum. *J Nippon Med Sch.* 2012;79(6):444-52. doi: 10.1272/jnms.79.444..
34. Im W, Chung JY, Kim SH, Kim M. Efficacy of autologous serum in human adipose-derived stem cells; cell markers, growth factors and differentiation. *Cell Mol Biol (Noisy-le-grand).* 2011 Mar 15;57 Suppl:OL1470-5. PMID: 21396339.
35. Bahn JJ, Chung JY, Im W, Kim M, Kim SH. Suitability of autologous serum for expanding rabbit adipose-derived stem cell populations. *J Vet Sci.* 2012 Dec;13(4):413-7. doi: 10.4142/jvs.2012.13.4.413.
36. de Paula AC, Zonari AA, Martins TM, Novikoff S, da Silva AR, Correlo VM, Reis RL, Gomes DA, Goes AM. Human serum is a suitable supplement for the osteogenic differentiation of human adipose-derived stem cells seeded on poly-3-hydroxybutyrate-co-3-hydroxyvalerate scaffolds. *Tissue Eng Part A.* 2013 Jan;19(1-2):277-89. doi: 10.1089/ten.TEA.2012.0189.
37. Koellensperger E, Bollinger N, Dexheimer V, Gramley F, Germann G, Leimer U. Choosing the right type of serum for different applications of human adipose tissue-derived stem cells: influence on proliferation and differentiation abilities. *Cytotherapy.* 2014 Jun;16(6):789-99. doi: 10.1016/j.jcyt.2014.01.007.
38. Tonarova P, Lochovska K, Pytlik R, Hubalek Kalbacova M. The Impact of Various Culture Conditions on Human Mesenchymal Stromal Cells Metabolism. *Stem Cells Int.* 2021 Mar 1;2021:6659244. doi: 10.1155/2021/6659244.
39. Dreher L, Elvers-Hornung S, Brinkmann I, Huck V, Henschler R, Gloe T, Klüter H, Bieback K. Cultivation in human serum reduces adipose tissue-derived mesenchymal stromal cell adhesion to laminin and endothelium and reduces capillary entrapment. *Stem Cells Dev.* 2013 Mar 1;22(5):791-803. doi: 10.1089/scd.2012.0051.
40. Hemedá H, Giebel B, Wagner W. Evaluation of human platelet lysate versus fetal bovine serum for culture of mesenchymal stromal cells. *Cytotherapy.* 2014 Feb;16(2):170-80. doi: 10.1016/j.jcyt.2013.11.004.
41. Piletz JE, Drivon J, Eisenga J, Buck W, Yen S, McLin M, Meruvia W, Amaral C, Brue K. Human Cells Grown With or Without Substitutes for Fetal

Human mesenchymal stem cells and human serum

Eur J Transl Myol 33 (1) 10834, 2023 doi: 10.4081/ejtm.2023.10834

Bovine Serum. Cell Med. 2018 Jun
6;10:2155179018755140.
doi:
10.1177/2155179018755140.

Disclaimer

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Submission: September 3, 2022

Revision received: November 30, 2022

Accepted for publication: November 30, 2022