

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/301340446>

The effect of oral N-acetylcysteine on serum creatinine in chronic kidney diseases patients under CABG surgery

Conference Paper · March 2016

CITATION

1

READS

12

4 authors:



[Mehdi deghani Firoozabadi](#)

Shahid Sadoughi University of Medical Scien...

20 PUBLICATIONS 11 CITATIONS

[SEE PROFILE](#)



[Ahmad Ebadi](#)

Ahvaz Jondishapour University of Medical Sc...

41 PUBLICATIONS 32 CITATIONS

[SEE PROFILE](#)



[Mohammad Ali Sheikhi](#)

Ahvaz Jondishapour University of Medical Sci...

20 PUBLICATIONS 9 CITATIONS

[SEE PROFILE](#)



[Hossein Rahmani](#)

Ahvaz Jondishapour University of Medical Sc...

39 PUBLICATIONS 11 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Systemic Inflammation in Stable COPD Patients due to Inhalation Toxicity [View project](#)

4th Iranian Cardiovascular Joint Congress

The effect of oral N-acetylcysteine on serum creatinine in chronic kidney diseases patients under CABG surgery

Mehdi Dehghani Firoozabadi 1, Ahmad Ebadi2*, Mohammad Ali Sheikhi 3, Hossein Rahmani 4,5

1. Department of Anesthesiology and Critical Care, Tehran Heart Center, Tehran University of Medical Sciences, Tehran, Iran
2. Department of Anesthesiology, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran *Corresponding author
3. Assistant Professor of Cardiovascular surgery MD, Department of Cardiac surgery, Atherosclerosis Research Center Golestan Hospital, Ahvaz Jundishapur University of Medical Sciences, Ahvaz
4. Toxicologist, Departments of Toxicology, Shahreza Branch, Islamic Azad University, Shahreza, Iran
5. Medical Research Center, Jundishapur Health Development Co, Tehran, Iran.

Abstract

Background: Many researches show risk factors of cardiac disease in individuals suffering from chronic kidney disease (CKD). In CKD patients who are candid CABG, there are great concerns about acute kidney injury. Increase in serum Creatinine is associated with increase in morbidity and mortality in CABG patients.

Objective: purpose of this study is investigating serum creatinine variation and GFR after oral NAC injection in CKD patients who are candid for CABG.

Material and Method: 50 patients candid for CABG were selected in blind design randomized, clinical trial in Golstan and Imam' hospitals in Ahvaz during 2010-2011. All patients had nephropathy type 2 based on kidney fund classification and had diagnostic criteria for chronic kidney diseases or it lasts for three months (GFR \leq 1.5). Intervention group received 600mg oral NAC twice a day from one day before surgery to 5 days after surgery through esophagus catheter and control group received distilled water with same amount and volume. Finally the serum creatinine, GFR and some other factors were measured.

Results: There is a significant difference in serum creatinine (P=0.002), GFR (P=0.002) and BUN (P=0.034) between control and intervention groups after 5 days.

Conclusion: Results of this study and its comparison with other studies showed great disagreement about effect of NAC on serum creatinine but considering lack of side effects or severe changes in serum creatinine in NAC receiving patients, it can be used as a safe drug.

Key words: N-acetylcysteine, serum creatinine, GFR, chronic kidney diseases (CKD), CABG


Introduction: Cardiac and kidney failure diseases are increasing everyday [1] and evidences show that failure in kidney function leads to cardiac failure and vice versa [2]. Many researches show risk factors of cardiac disease in individuals suffering from chronic kidney disease (CKD) [3-4] and it is observed that kidney diseases increase cardiovascular diseases, mortalities and morbidities [5] such that we can say patients suffering from CKD will die as a result of cardiovascular diseases not advanced kidney failures [6] because rate of stroke in CKD patients is high and can lead to death related to cardiovascular diseases [7] and it is estimated that 13% of CKD patients suffer from myocardial stroke [8-9]. In CKD patients who are candid for coronary artery bypass grafting (CABG), there are great concerns about acute kidney failure and rate of acute kidney injury (AKI) after cardiac surgery varies from 1-3% dependent on patients' characteristics [10]. Various studies have reported that increase in serum Creatinine is associated with increase in morbidity and mortality in CABG patients under pump [11] and slight increase in serum Creatinine leads to cardiovascular injuries and stroke [12]. Reports indicated that increase in serum creatinine is seen in 20-40% of patients hospitalized because of cardiac failure [13-15]. Creatinine is used for estimating glomerular filtration rate (GFR) [16] and one of risk factors in cardiovascular diseases in CKD patients is reducing GFR after increasing serum creatinine [17]. Results of researchers show that there is difference about dependence or independence of serum creatinine variations relative to GFR [18-21] because serum creatinine is influenced by factors like age, gender, race and body mass index and it is necessary to consider these important factors to understand relationship between increasing Creatinine and reducing GFR with accurate information and development of cardiovascular diseases [22]. Rep [27]. It is necessary to mention that one of important mechanisms for kidney injury is oxidative stress [28] and NAC is a sulfhydryl-containing compound which has strong anti-oxidative features [29]. NAC is suitable for cardiac function improvement after myocardial stroke [30]. Studies have stated that NAC can reduce serum creatinine [31] but instead of these studies NAC has a contrasted history about preventing kidney injuries [32-33] and even reduction in serum creatinine after NAC injection was debated by researchers [34-35]. Therefore, purpose of this study is investigating serum creatinine variation and GFR after oral NAC injection in CKD patients who are candid for CABG.

Material and Method:


after approval of ethic committee in by Ahvaz Jundishapur University of Medical Sciences and consent form signed by participants, 50 patients candid for CABG were selected in blind design randomized, clinical trial in Golstan and Imam' hospitals in Ahvaz during 2010-2011. All patients had nephropathy type 2 based on kidney fund classification and had diagnostic criteria for chronic kidney diseases or it lasts for three months (GFR \leq 90, Cr $>$ 1.5). Patients who had dialysis before surgery or emergency operation or kidney implant or received contrast drug one week before study were excluded from the study. Other patients were assigned randomly to groups receiving NAC and placebo. Intervention group received 600mg oral NAC twice a day from one day before surgery to 5 days after surgery through esophagus catheter and control group received distilled water with same amount and volume. Patients received 0.1mg/kg intramuscular morphine and 25mg promethazine. In the case of hemodynamic stability and pain control patient was removed from mechanical ventilation. Hemodynamic management in patients includes keeping MAP $>$ 65 with crystalloid and Clويد-infusion Dopamine and epinephrine were injected, when necessary

Results: results of study showed that primary characteristic of patients in study and their histories in both control and intervention group had not significant differences with each other (table 1). Data analysis indicated that cardiac EF of both groups had no significant differences (P=0.164). Mean total bypass time was 72.72 \pm 16.35 for control group and 88.40 \pm 12.88 for intervention group. This time had significant differences between both groups (P=0.003). from BUN concentration comparison we can show that using NAC will significantly decrease BUN in CABG surgery (P=0.034). This reduction in concentration in intervention group is higher than control group. Comparing Na, K, HCO₃ concentration between both control and intervention groups in different times showed insignificant differences between both groups and we can conclude that using NAC has not significant effect on Na, K and HCO₃ concentration in CABG patients. Based on Friedman analysis there was a significant statistical differences in creatinine variation one day before surgery to 5 days after surgery in control group such that mean creatinine was increased from 1.60 mg/dl to 1.90 mg/dl in day 0. Based on this test, there was a significant difference in creatinine variation from one day before surgery and 5 days after surgery such that its mean reduced from 1.80 mg/dl to 1.40 mg/dl. Based on Mann-Whitney test there is a significant difference between creatinine in day 5 between control and intervention groups (P=0.002). Mean creatinine was increased in control group but decreased in intervention group (figure 1). Based on this test, there is a significant difference between groups from day 5. Based on Friedman test, there is a significant statistical difference between GFR from one day before surgery to 5 days after surgery in control group such that mean GFR was reduced from 47.03 in day 0 to 41.62 ml in minutes. Based on this test there was a significant statistical difference between GFR from day 1 before surgery to 5 days after surgery in intervention group such that median GFR increased from 40.70 in day 0 to 51.90 ml in minutes

Mean GFR



Mean creatinine



Discussion and Conclusion:

CKD patients cannot participate in clinical trials because there are great concerns about side effects of drugs and methods but it is clear that there is a need for more studies to increase life quality of these patients. As results of this study show, changes in serum creatinine has increased in control group but decreased in intervention group and caused GFR difference between both groups such that mean GFR increased in intervention group but decreased in control group. Adabag(36) et.al, by studying effect of oral NAC on AKI prevention after cardiac surgery, had reported that serum creatinine in NAC group has increases 0.45 \pm 0.7 mg/dl but 0.55 \pm 0.9mg/dl in control group, but this difference is not significant and serum creatinine increase in NAC group in this study was lower than control group (P=0.53). Staniloae(37) et.al in a study on NAC effects on preventing nephropathy caused by contrast in angiography patients with kidney failures showed that serum creatinine was increased in 4.2% intervention group patients while in control group 6.5% patients had increase in serum creatinine but this was not significant (P=0.38). But like our study increase in serum creatinine was lower in intervention group. Following otheon kidney failure in high risk CABG patients and showed that in 29.7% of NAC receiving group serum creatinine was increased but in control group 29% had increase which was not significant (p=0.89). Webb(40) et.al, by studying effect of NAC on preventing nephropathy caused by contrast in cardiac catheterization, showed that in NAC group 7.3% patients had increase in serum creatinine while this increase occurred in 5.7% patients in control group and in their study, NAC has increased serum creatinine but was not statistically significant (p=0.57) and contrasted with our results. It is necessary to mention that in this study we have investigated effect of NAC on other factors like HCO₃, K, Na and BUN which had significant difference in BUN. Therefore, it is suggested that other studies investigate creatinine variation and other factors like BUN.

References

- 1-Metra M, Cotter G, Gheorghiade M, Dei Cas L, Voors AA. The role of the kidney in heart failure. Eur Heart J. 2012;33:2135-42
- 2-Lekawanvijit S, Kompa AR, Wang BH, Kelly DJ, Krum H. Cardiorenal syndrome: the emerging role of protein-bound uremic toxins. Circ Res. 2012;111: 1470-1483
- 3-Murnner P, Hamm LL, Kusek JW, Chen J, Whelton PK, He J. The prevalence of nontraditional risk factors for coronary heart disease in patients with chronic kidney disease. Ann Intern Med. 2004; 140: 94-107
- 4-Sarnak MJ, Coronado BE, Greene T, Wang SR, Kusek JW, Beck GJ, Levey AS. Cardiovascular disease risk factors in chronic renal insufficiency. Clin Nephrol. 2002; 57: 327-335
- 5-Tonelli M, Wiebe N, Cullerton B et al. Chronic kidney disease and mortality risk: a systematic review. J Am Soc Nephrol. 2006; 17: 2034-2047
- 6-Sarnak MJ, Levey AS, Schoolwerth AC, Corsh J, Cullerton B, Hamm LL, McCullough PA, Kasike BL, Kelepouris E, Klag MJ, Parfrey P, Pfeffer M, Raij L, Spinosa DJ, Wilson PW. Kidney disease as a risk factor for development of cardiovascular disease: a statement from the American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. Hypertension. 2003;42:1050-1065
- 7-Meisinger C, Doring A, Lowell H. Chronic kidney disease and risk of incident myocardial infarction and all-cause and cardiovascular disease mortality in middle-aged men and women from the general population. Eur Heart J. 2006; 27: 1245-1250
- 8- Bansal N. Clinically silent myocardial infarctions in the CKD community. Nephrology Dialysis Transplantation. 2012; 27(9): 3387-3391
- 9-Risk DV, Gutierrez O, Levin EB et al. Prevalence and prognosis of unrecognized myocardial infarctions in chronic kidney disease. Nephrol Dial Transplant. 2011. [Epub ahead of print December 13]
- 10-Mangan, Om Diamondstone LS Ramsay JC, et al. multicenter study of preoperative ischemia Research Group Renal dysfunction after my cordiel revascularisations is at el Ann Intern Med. 1998; 128:194-203
- 11-Lee SY, Choi JB, Lee MK. Changes of Renal Function and Treatment after CABG in Patients with Elevated Serum Creatinine. Korean J Thorac Cardiovasc Surg. 2005; 38(2):146-151.
- 12-Hoke M, Amighi J, Mlekusch W, Schlager C, Exner M, Sabetti S, Dick P. Cystatin C and the risk for cardiovascular events in patients with asymptomatic carotid atherosclerosis. Stroke. 2010;41:674-679
- 13-Butler J, Chrosvsky D, Prastak H, McNeill A, Cody R. Renal function, health outcomes, and resource utilization in acute heart failure: a systematic review. Circ Heart Fail. 2010;3:726-745.
- 14-Metra M, Nadari S, Parrinello G, Bordonali T, Bugatti S, Danesi R, Fontanella B, Lombardi C, Milani P, Verzura G, Cotter G, Dittich H, Massie BM, Dei Cas L. Worsening renal function in patients hospitalised for acute heart failure: clinical implications and prognostic significance. Eur J Heart Fail. 2008;10:188-195.